



Semileptonic decays and tests of lepton flavour universality at Belle II

Christoph Schwanda
Austrian Academy of Sciences
Representing the collaboration


21st Lomonosov Conference on Elementary Particle Physics
August 24-30, 2023, Moscow State University



Search for LFU violation

Motivation

- Physics beyond the Standard Model can spoil the universality of lepton flavour couplings
 - Probing lepton flavour universality (LFU) is thus a promising, theoretically clean avenue for searching for New Physics
- Semileptonic B meson decays combine high rate and low experimental background
 - They thus present an excellent tool for LFU searches
 - There is a long-standing 3σ LFU anomaly in semitauonic B decays



Outline

Belle II results covered in this presentation

- Measurement of $R(D^*)$
[189/fb, Lepton Photon 2023]

$$R(D^*) = \frac{\mathcal{B}(B \rightarrow D^* \tau \nu_\tau)}{\mathcal{B}(B \rightarrow D^* \ell \nu_\ell)}$$

- Measurement of $R(X)$
[189/fb, EPS-HEP 2023]

$$R(X) = \frac{\mathcal{B}(B \rightarrow X \tau \nu_\tau)}{\mathcal{B}(B \rightarrow X \ell \nu_\ell)}$$

- Tests of light-lepton universality in angular asymmetries of $B \rightarrow D^* \ell \nu$
[189/fb, [arXiv:2308.02023](#), submitted to Phys. Rev. Lett.]
- Test of light-lepton universality in inclusive semileptonic B meson decays
[189/fb, [arXiv:2301.08266](#), [Phys. Rev. Lett. 131, 051804 \(2023\)](#)]

The Belle II detector



KEK
Tsukuba, Japan

Vertex detector
2 layers of DEPFET pixels (PXD) and
4 layers of silicon strips (SVD)
Vertex resolution $\sim 15\mu\text{m}$

Central drift chamber
Spatial resolution $\sim 100\mu\text{m}$
 dE/dx resolution: 5%
 p_T resolution: 0.4%

KLM
Instrumented flux return

Electromagnetic Calorimeter
Energy resolution: 1.6 - 4%


$E_{\text{cm}} = 10.58 \text{ GeV}$
($\Upsilon(4S)$ resonance)

$4 \text{ GeV } e^+$

Forward and barrel Part. Id.
K eff. 90%, fake π rate 5%

Belle II timeline

Luminosity projection






Measurement of $R(X)$


[189/fb, EPS-HEP 2023]

Reconstruction

- Reconstruct one B meson in a hadronic decay mode (B_{tag})
- Reconstruct a leptonic τ within remaining particles ($\tau \rightarrow e\nu\nu, \mu\nu\nu$)
 - $p_{T,\text{lab}}(e) > 0.3/0.5 \text{ GeV}, p_{T,\text{lab}}(\mu) > 0.4/0.7 \text{ GeV}$
- The remaining particles on the signal side are collectively referred to as X
- Main challenge: correct model of backgrounds
 - Data-driven $X\ell\nu$ re-shaping using the M_X distribution in the $p_\ell^B > 1.4 \text{ GeV}$ region



Signal extraction



- From the p_ℓ^B vs. M_{miss}^2 distribution, separately for e and μ events
- 34 bins in p_ℓ^B vs. M_{miss}^2
- 2×4 fit components: $X\tau\nu$, $X\ell\nu$, $B\bar{B}$ background (fakes and secondaries), continuum (off-resonance data, yield constrained)

Results

preliminary



$$R(X) = \frac{\mathcal{B}(B \rightarrow X\tau\nu_\tau)}{\mathcal{B}(B \rightarrow X\ell\nu_\ell)}$$

$$R(X_{\tau/e}) = 0.232 \pm 0.042, [0.020 \text{ (stat)}, 0.037 \text{ (syst)}]$$

$$R(X_{\tau/\mu}) = 0.222 \pm 0.057, [0.027 \text{ (stat)}, 0.050 \text{ (syst)}]$$

$$R(X_{\tau/\ell}) = 0.228 \pm 0.039, [0.016 \text{ (stat)}, 0.036 \text{ (syst)}]$$

- Total uncertainty: 17.5%
- Largest systematics:
signal and background model
- Consistent with previous measurements
from LEP, the SM expectation and
constraints from $R(D^{(*)})$






Measurement of $R(D^*)$

[189/fb, Lepton Photon 2023]




Reconstruction



- Reconstruct one B meson in a hadronic decay mode (B_{tag})
- Reconstruct a D^* and a leptonic τ decay ($\tau \rightarrow e\nu\nu, \mu\nu\nu$) on the signal-side within remaining particles
 - Three D^* modes:
 $D^{*+} \rightarrow D^0\pi^+, D^+\pi^0, D^{*0} \rightarrow D^0\pi^0$
- Rest of the event: no charged tracks, no π^0 candidates
- Main challenge: three neutrinos in the final state, significant, sometimes poorly understood ($D^{**}\ell\nu$) backgrounds


Data-driven validation of the background model

Using different side bands



$R(D^*)$ signal extraction

- Two-dimensional binned likelihood fit to
 - E_{ECL} : energy remaining in the calorimeter after removing all reconstructed particles
 - $M_{\text{miss}}^2 = (p_{e^+e^-} - p_{B_{\text{tag}}} - p_{D^*} - p_{\ell})^2$: missing mass of the event






Results

zoomed

$D^*\tau\nu$ enhanced

$$1.5 < M_{\text{miss}}^2 < 6.0 \text{ (GeV}/c^2)^2$$



$$R(D^*) = \frac{\mathcal{B}(B \rightarrow D^* \tau \nu_\tau)}{\mathcal{B}(B \rightarrow D^* \ell \nu_\ell)}$$

$$R(D^*) = 0.267^{+0.041}_{-0.039}(\text{stat.})^{+0.028}_{-0.033}(\text{syst.})$$

- First $R(D^*)$ result from Belle II data
- Main systematics: MC statistics, shape of E_{ECL}
- Consistent both with the SM and other experimental determinations of $R(D^*)$

preliminary

Tests of light-lepton universality in angular asymmetries of $B \rightarrow D^* \ell \nu$

[189/fb, [arXiv:2308.02023](#), submitted to Phys. Rev. Lett.]

Definition of angular observables in $B \rightarrow D^* \ell \nu$

$$\mathcal{A}_x(w) \equiv \left(\frac{d\Gamma}{dw} \right)^{-1} \left[\int_0^1 - \int_{-1}^0 \right] dx \frac{d^2\Gamma}{dw dx}$$

with $x = \cos \theta_\ell$ for A_{FB}

$\cos 2\chi$ for S_3


$\cos \chi \cos \theta_V$ for S_5

$\sin \chi \cos \theta_V$ for S_7

$\sin 2\chi$ for S_9

$$\Delta \mathcal{A}_x(w) \equiv \mathcal{A}_x^\mu(w) - \mathcal{A}_x^e(w)$$

$$w \equiv \frac{m_{B^0}^2 + m_{D^*}^2 - q^2}{2m_B m_{D^*}}$$



in W rest frame



- Theory motivation for the study of these observables:
Phys. Rev. D 107 (2023) 1, 015011
[arXiv:2206.11283 \[hep-ph\]](https://arxiv.org/abs/2206.11283)

Angular asymmetries in $B \rightarrow D^* \ell \nu$

- Reconstruction

- $B^0 \rightarrow D^{*-} \ell^+ \nu_\ell$ decays are reconstructed in hadronically tagged $\Upsilon(4S)$ events
- Signal is extracted from the M_{miss}^2 distributions
- Asymmetries are extracted in the full w region ($w_{\text{incl.}}$), in w_{low} ($1 < w < 1.275$) and in w_{high} ($1.275 < w < 1.503$)

- Result



preliminary

Test of light-lepton universality in inclusive semileptonic B meson decays


[189/fb, [arXiv:2301.08266](#), [Phys. Rev. Lett. 131, 051804 \(2023\)](#)]

$$R(X_{e/\mu})$$

- Analysis and background correction technique is shared with the $R(X)$ measurement
- The ratio of inclusive semileptonic decays to e and to μ is obtained in the region $p_{\ell}^B > 1.3$ GeV
- Result

$$R(X_{e/\mu}) = 1.007 \pm 0.0009(\text{stat}) \pm 0.019(\text{syst})$$



$$R(X_{e/\mu}) = \frac{\mathcal{B}(B \rightarrow Xe\nu_e)}{\mathcal{B}(B \rightarrow X\mu\nu_\mu)}$$



- Main systematics: lepton identification
- Most precise LFU test so far (2%), consistent with SM


Summary and conclusion

- Belle II has probed lepton flavour universality (LFU) in semileptonic B meson decays
 - First measurement of inclusive semitauonic B decays at the $\Upsilon(4S) - R(X)$
 - First measurement of $B \rightarrow D^* \tau \nu$ at Belle II – $R(D^*)$
 - Forward-backward asymmetry (and other angular observables) in $B \rightarrow D^* \ell \nu$ separately for $\ell = e, \mu$
 - Most precise test of light lepton universality in inclusive semileptonic B decays
- So far, results are consistent with the SM and previous experimental findings. There is still large room for improvement as more Belle II data is collected



Backup




Untagged vs. Tagged



Hadronic tagging at Belle II



Comput Softw Big Sci (2019) 3: 6.



$$M_{bc} = \sqrt{E_{\text{beam}}^2/4 - (p_{B_{\text{tag}}}^{\text{cm}})^2} > 5.27 \text{ GeV}/c^2$$

- The hadronic FEI employs over 200 boosted decision trees to reconstruct 10000 B decay chains
 - $\epsilon_{B^+} \approx 0.5 \%$, $\epsilon_{B^0} \approx 0.3 \%$ at low purity (about 50% increase with respect to the Belle tag)