




status and prospects



BEAUTY
2019


18th INTERNATIONAL CONFERENCE
ON B-PHYSICS AT FRONTIER MACHINES
Ljubljana, Slovenia
September 30 - October 4, 2019

Gagan Mohanty



☞ MC simulations yet to include embedded random triggers to correctly represent the beam background effects and electronic noise

First-generation e^+e^- flavor factories




Success culminated in 2008 Nobel prize in Physics

Rich legacy left for next-gen expt. EPJ C74, 3026 (2014)



So, why another e^+e^- flavor factory?

- ❑ Precision CKM metrology → Standard Model (SM) candle
- ❑ New CP violating phase? → CP violation in B and D decays
- ❑ Any imprint of new physics beyond SM in FCNC transitions? → radiative and electroweak penguin decays
- ❑ How about charged Higgs boson or leptoquark? → tree-level B decay to $\tau\nu$ or $D^{(*)}\tau\nu$ final state
- ❑ New physics in tau lepton sector → search for lepton flavor violating (LFV) tau decays
- ❑ Can we chase down dark matter from bottom? → hidden dark sector



@ SuperKEKB will address these questions with almost two orders of magnitude larger dataset than Belle+BABAR

Snapshots of what can achieve?

Observables	Expected the. accuracy	Expected exp. uncertainty	Facility (2025)
UT angles & sides			
ϕ_1 [°]	***	0.4	Belle II
ϕ_2 [°]	**	1.0	Belle II
ϕ_3 [°]	***	1.0	LHCb/Belle II
$ V_{cb} $ incl.	***	1%	Belle II
$ V_{cb} $ excl.	***	1.5%	Belle II
$ V_{ub} $ incl.	**	3%	Belle II
$ V_{ub} $ excl.	**	2%	Belle II/LHCb
CP Violation			
$S(B \rightarrow \phi K^0)$	***	0.02	Belle II
$S(B \rightarrow \eta' K^0)$	***	0.01	Belle II
$\mathcal{A}(B \rightarrow K^0 \pi^0) [10^{-2}]$	***	4	Belle II
$\mathcal{A}(B \rightarrow K^+ \pi^-) [10^{-2}]$	***	0.20	LHCb/Belle II
(Semi-)leptonic			
$\mathcal{B}(B \rightarrow \tau \nu) [10^{-6}]$	**	3%	Belle II
$\mathcal{B}(B \rightarrow \mu \nu) [10^{-6}]$	**	7%	Belle II
$R(B \rightarrow D \tau \nu)$	***	3%	Belle II
$R(B \rightarrow D^* \tau \nu)$	***	2%	Belle II/LHCb
Radiative & EW Penguins			
$\mathcal{B}(B \rightarrow X_s \gamma)$	**	4%	Belle II
$A_{CP}(B \rightarrow X_{s,d} \gamma) [10^{-2}]$	***	0.005	Belle II
$S(B \rightarrow K_S^0 \pi^0 \gamma)$	***	0.03	Belle II
$S(B \rightarrow \rho \gamma)$	**	0.07	Belle II
$\mathcal{B}(B_s \rightarrow \gamma \gamma) [10^{-6}]$	**	0.3	Belle II
$\mathcal{B}(B \rightarrow K^* \nu \bar{\nu}) [10^{-6}]$	***	15%	Belle II
$R(B \rightarrow K^* \ell \ell)$	***	0.03	Belle II/LHCb
Charm			
$\mathcal{B}(D_s \rightarrow \mu \nu)$	***	0.9%	Belle II
$\mathcal{B}(D_s \rightarrow \tau \nu)$	***	2%	Belle II
$A_{CP}(D^0 \rightarrow K_S^0 \pi^0) [10^{-2}]$	**	0.03	Belle II
$ q/p (D^0 \rightarrow K_S^0 \pi^+ \pi^-)$	***	0.03	Belle II
$A_{CP}(D^+ \rightarrow \pi^+ \pi^0) [10^{-2}]$	**	0.17	Belle II
Tau			
$\tau \rightarrow \mu \gamma [10^{-10}]$	***	< 50	Belle II
$\tau \rightarrow e \gamma [10^{-10}]$	***	< 100	Belle II
$\tau \rightarrow \mu \mu \mu [10^{-10}]$	***	< 3	Belle II/LHCb

👉 From Belle II physics book [arXiv:1808.10567](https://arxiv.org/abs/1808.10567)

Precision CKM metrology

Direct and mixing-induced CP violation in B decays

(Semi-)leptonic B decays

Radiative & electroweak penguins




Vibrant charm program

Search of LFV tau decays



New intensity frontier machine


- Targets to deliver e^+e^- collisions at a peak luminosity of $8 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$, 40 times that of KEKB
 - Increase beam currents **twice**
 - Reduce beam size by **20 times**



➤ First new particle collider after LHC!


How far have we gone?

Belle II online luminosity Exp: 7-8 - All runs




- Phase 2 (2018): beam commissioning (establish nano-beam scheme, reach the KEKB luminosity, and measure beam backgrounds) as well as for doing some physics with partial vertex detector
- Phase 3 (2019 onward): physics run with almost complete vertex detector

- Reached $\beta_y^* = 33$ mm in 2018
- Went down $\beta_y^* = 2$ mm by end of Summer 2019 (with Belle II off) → starting point for fall run
- Design luminosity requires one more order-of-magnitude jump to $\beta_y^* = 0.3$ mm



- Currents achieved: 880 (940) mA for e⁺ (e⁻) beam → need 3 (4)× scale up




- ❑ Close to 950 researchers from 112 institutions in 26 countries
- ❑ Slovenia is an important player
 - 👉 Leadership position, as well as key contributions to detector, reconstruction software and computing



: A 21st century HEP experiment

➔ Designed to operate with a performance similar or better than Belle, but in a harsh beam background condition



Two detector highlights

Barrel PID (imaging TOP): Japan, US, Slovenia and Italy


👉 Example of Cherenkov-photon paths for 2 GeV pion and kaon traversing in a TOP quartz bar

Micro-channel-plate (MCP)
PMTs; 512 channels; 50 ps
resolution


Incoming
 π/K track

Cherenkov angle:
 $\cos \theta_c = 1/n\beta$

Photon from π^+
Photon from K^+





VXD (6 layer Si for vertexing & inner tracking)



- Beam-pipe $r = 10$ mm
 - DEPFET pixels: Germany, Czech Republic, Spain...
 - Layer 1 $r = 14$ mm
 - Layer 2 $r = 22$ mm (1/6 now, rest in 2020)
 - DSSD (double sided micro-strips)
 - Layer 3 $r = 38$ mm (Australia)
 - Layer 4 $r = 80$ mm (India)
 - Layer 5 $r = 115$ mm (Austria)
 - Layer 6 $r = 140$ mm (Japan)
- FWD/BWD
Italy

A performance example: dE/dx in CDC




- ❑ Obtained with early calibrations in the hadronic event sample
- ❑ Important role in identifying charged particles
- 👉 More performance results will be in Tenchini's talk


A physics example: fresh from




Belle II preliminary $\int \mathcal{L} dt = 5.15 \text{ fb}^{-1}$




Belle II preliminary $\int \mathcal{L} dt = 5.15 \text{ fb}^{-1}$



Belle II preliminary $\int \mathcal{L} dt = 5.15 \text{ fb}^{-1}$




Belle II preliminary $\int \mathcal{L} dt = 5.15 \text{ fb}^{-1}$



- Study of untagged $B \rightarrow D^* \ell \nu_\ell$ channel with 5.15 fb^{-1} early phase-3 data
- Over 1k signal events found in both electron and muon channels

☞ For more on physics, see talks by Yonenaga (rare decays), Graziani (dark sector), Yusa (CP violation in B decays), and Kwon (charm)

Prospects: physics harvesting



All the details are in
 "The Belle II Physics Book"
[arXiv:1808.10567](https://arxiv.org/abs/1808.10567)

👉 Adapted from Forti's talk at EPS-HEP 2019

Prospects: improvements to detector

❑ Short term:


- Replacement of conventional MCP-PMTs with Atomic-layer-deposition (ALD) PMTs for the TOP counters
- Complete installation of PXD layer-2
- DAQ upgrade

❑ Medium term:

- Looking at options for making the detector more resilient against beam-induced background and radiation bursts

❑ Longer term:

- Started to think about possibilities for luminosity upgrade; e.g., Belle II VXD open workshop <http://indico.cern.ch/event/810687/>



Closing words

- ❑ Belle II will probe new physics at the intensity frontier → complementary to high- p_T programs of ATLAS and CMS
- ❑ As for LHCb, there is healthy competition and complementarity between the two experiments
- ❑ 1st physics run in Spring 2019 has completed delivering $\sim 6.5 \text{ fb}^{-1}$ → fall run is about to begin
- ❑ Detector and machine initial performances have been good; we expect the road ahead to be bit long before achieving the design goal




Additional information

Comparison: KEKB vs. SuperKEKB

parameters		KEKB		SuperKEKB		units
		LER	HER	LER	HER	
Beam energy	E_b	3.5	8	4	7	GeV
Half crossing angle	ϕ	11		41.5		mrad
Horizontal emittance	ϵ_x	18	24	3.2	4.6	nm
Emittance ratio	κ	0.88	0.66	0.37	0.40	%
Beta functions at IP	β_x^*/β_y^*	1200/5.9		32/0.27	25/0.30	mm
Beam currents	I_b	1.64	1.19	3.60	2.60	A
beam-beam parameter	ξ_y	0.129	0.090	0.0881	0.0807	
Luminosity	L	2.1×10^{34}		8×10^{35}		$\text{cm}^{-2}\text{s}^{-1}$

Beam backgrounds



- e^+e^- colliders are clean, however at high L_{peak} value beam backgrounds can become a challenge
- At the highest luminosities, QED processes e.g., $e^+e^- \rightarrow e^+e^-(\gamma)$ & $e^+e^- \rightarrow e^+e^-e^+e^-$ will dominate

- Currently, single beam backgrounds are dominant, larger for the e^+ beam
 - beam-gas (residual gas in beam-pipe)
 - Touschek (intra-bunch scattering)
 - injection-induced
 - “dust events” (occasional large losses)
- CDC HV trips with large background
- Beam abort protection against spikes due to radiation
- Simulation and collimator studies

