

Charm Physics at Belle and Belle II experiments

Longke LI (李龙科)

On behalf of the Belle and Belle II Collaborations

lilk@ucmail.uc.edu

University of Cincinnati (UC)



the 2024 International Workshop on Future Tau Charm Facilities (FTCF 2024)
Jan 18, 2024 @Univ. of Sci & Tech of China (my alma mater ♡)

Outline




- 1 Charm sample at Belle and Belle II
- 2 Charm lifetime measurements
- 3 Branching fraction and decay asymmetry parameter
- 4 Search for rare or forbidden decay
- 5 Charm CP violation searches
- 6 Summary

Outline







- 1 Charm sample at Belle and Belle II
- 2 Charm lifetime measurements
- 3 Branching fraction and decay asymmetry parameter
- 4 Search for rare or forbidden decay
- 5 Charm CP violation searches
- 6 Summary

Charm production at Belle and Belle II

- At Belle (II), e^+e^- mainly collide at 10.58 GeV to make $\Upsilon(4S)$ resonance decaying into $B\bar{B}$ in 96% of the time.
- Meanwhile, continuum processes $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) have large cross sections.
- Two ways to produce the charm sample: $e^+e^- \rightarrow c\bar{c}$ ($\sigma = 1.3$ nb), and $B \rightarrow$ charm decays.



Comparison of available charm samples from Charm factories, B-factories, hadron colliders

Experiment	Machine	Operation	C.M.	Luminosity	N_{prod}	Efficiency	Characters
	(e^+e^-)	2003-2008	3.77 4.18 GeV	0.8 fb ⁻¹ 0.6 fb ⁻¹	$D^{0,+}$: $\mathcal{O}(10^6)$ D_s^+ : 6×10^5	~10-30%	<ul style="list-style-type: none"> ☺ extremely clean environment ☺ pure D-beam, almost no bkg ☺ quantum coherence ☹ no CM boost, no time-dept analyses
		2010-2011 (2021-)	3.77 GeV	2.9 (8 → 20) fb ⁻¹	$D^{0,+}$: $10^7 (\rightarrow 10^8)$		
	(e^+e^-)	2016-2019	4.18-4.23 GeV	7.3 fb ⁻¹	D_s^+ : 5×10^6	★★★	
		2014+2020	4.6-4.7 GeV	4.5 fb ⁻¹	Λ_c^+ : 0.8×10^6 ★☆☆		
	SuperKEKB (e^+e^-)	2019-	10.58 GeV	0.4 (→ 50) ab ⁻¹	D^0 : $6 \times 10^8 (\rightarrow 10^{11})$ $D_{(s)}^+$: $10^8 (\rightarrow 10^{10})$ Λ_c^+ : $10^7 (\rightarrow 10^9)$	$\mathcal{O}(1-10\%)$	<ul style="list-style-type: none"> ☺ clear event environment ☺ high trigger efficiency ☺ high-efficiency detection of neutrals ☺ time-dependent analysis ☹ smaller cross-section than hadron colliders
	KEKB (e^+e^-)	1999-2010	10.58 GeV	1 ab ⁻¹	D^0 : 10^9 $D_{(s)}^+$: 10^9 Λ_c^+ : 10^8		
	PEP-II (e^+e^-)	1999-2008	10.58 GeV	0.5 ab ⁻¹	6×10^8 ★★☆	★★	
	Tevatron $(p\bar{p})$	2002-2011	1.96 TeV	9.6 fb ⁻¹	1.3×10^{11}	$\mathcal{O}(0.1\%)$	<ul style="list-style-type: none"> ☺ very large production cross-section ☺ large boost ☺ excellent time resolution ☹ dedicated trigger required
	LHC (pp)	2011,2012 2015-2018 (2022-2025,2029-)	7+8 TeV 13 TeV	1+2 fb ⁻¹ 6 fb ⁻¹ (→ 23 → 50) fb ⁻¹	5×10^{12} 10^{13} ★★★★★		

Here uses $\sigma(D^0 \bar{D}^0 @ 3.77 \text{ GeV}) = 3.61 \text{ nb}$, $\sigma(D^+ D^- @ 3.77 \text{ GeV}) = 2.88 \text{ nb}$, $\sigma(D_s^* D_s @ 4.17 \text{ GeV}) = 0.967 \text{ nb}$; $\sigma(c\bar{c} @ 10.58 \text{ GeV}) = 1.3 \text{ nb}$ where each $c\bar{c}$ event averagely has 1.1/0.6/0.3 $D^0/D^+/D_s^+$ yields; $\sigma(D^0 @ \text{CDF}) = 13.3 \mu\text{b}$, and $\sigma(D^0 @ \text{LHCb}) = 1661 \mu\text{b}$, mainly from *Int. J. Mod. Phys. A* **29**(2014)24,14300518.

Outline



- 1 Charm sample at Belle and Belle II
- 2 Charm lifetime measurements**
- 3 Branching fraction and decay asymmetry parameter
- 4 Search for rare or forbidden decay
- 5 Charm CP violation searches
- 6 Summary



Charm lifetimes at Belle II

PRL 127, 211801 (2021), PRL 130, 071802 (2023), PRD 107, L031103 (2023)

- Hadron lifetimes are difficult to calculate theoretically, as they depend on nonperturbative effects arising from quantum chromodynamics (QCD).
- Comparing calculated values with measured values improves our understanding of QCD. [(FLAG) EPJC 82, 869 (2022)]
- At Belle II, the decay-time resolution is $\times 2$ better than that at Belle/BABAR.
- Based on the early Belle II dataset, the most precise charm lifetimes are measured: $\tau(D^0) = 410.5 \pm 1.1 \pm 0.8$ fs, $\tau(D^+) = 1030.4 \pm 4.7 \pm 3.1$ fs, and $\tau(\Lambda_c^+) = 203.20 \pm 0.89 \pm 0.77$ fs as first precision measurements at Belle II.



Precise measurement of D_s^+ lifetime at Belle II




PRL 131, 171803 (2023)

- A clear sample of $D_s^+ \rightarrow \phi\pi^+$ with 116K signals and 92% purity, is obtained using 207 fb^{-1} .
- Lifetime determined from unbinned ML fit to lifetime (t). The likelihood function for i th event:






$$\mathcal{L}(\tau | t^i, \sigma_t^i) = f_{\text{sig}} P_{\text{sig}}(t^i | \tau, \sigma_t^i) P_{\text{sig}}(\sigma_t^i) + (1 - f_{\text{sig}}) P_{\text{bkg}}(t^i | \tau, \sigma_t^i) P_{\text{bkg}}(\sigma_t^i)$$

where $P_{\text{sig}}(\sigma_t^i)$ and $P_{\text{bkg}}(\sigma_t^i)$ exist to avoid the Punzi bias [arXiv:physics/0401045].

- Result: $\tau_{D_s^+} = (499.5 \pm 1.7 \pm 0.9) \text{ fs}$; the world most precise measurement to date.
- The small systematic uncertainty demonstrates the excellent performance and understanding of the Belle II detector.



Charm lifetime summary




- In all cases except for Ω_c^0 , Belle II has made the world's highest precision measurement (in some cases after 20 years)
- For Ω_c^0 , the Belle II measurement confirms the longer lifetime measured by LHCb.

Outline

- 1 Charm sample at Belle and Belle II
- 2 Charm lifetime measurements
- 3 Branching fraction and decay asymmetry parameter**
- 4 Search for rare or forbidden decay
- 5 Charm CP violation searches
- 6 Summary

Branching fraction of charmed meson Cabibbo-suppressed decays

- Cabibbo-suppressed (CS) hadronic decays of charm mesons provide a powerful means to search for new physics. It is important to measure such decays with high precision.
- The singly Cabibbo-suppressed (SCS) charm decays are essential probes of CP violation in the charm sector and new physics beyond the standard model.
- The large inclusive samples of charm hadrons at Belle and Belle II provide a good platform to measure the relative branching fractions of charmed hadron decays.
- Based on Belle entire dataset (980 ab^{-1}), the first or most precise \mathcal{B} results for charmed meson decays were reported:
- SCS decay:
 - $\mathcal{B}(D^+ \rightarrow K^+ K^- \pi^+ \pi^0) = (7.08 \pm 0.08 \pm 0.16 \pm 0.20) \times 10^{-3}$
 - $\mathcal{B}(D_s^+ \rightarrow K^+ \pi^- \pi^+ \pi^0) = (9.44 \pm 0.34 \pm 0.28 \pm 0.32) \times 10^{-3}$
 - $\mathcal{B}(D_s^+ \rightarrow K^+ K^- K_S^0 \pi^+) = (1.29 \pm 0.14 \pm 0.04 \pm 0.11) \times 10^{-4}$
- DCS decay:
 - $\mathcal{B}(D^+ \rightarrow K^+ \pi^- \pi^+ \pi^0) = (1.05 \pm 0.07 \pm 0.02 \pm 0.03) \times 10^{-3}$
(confirm BESIII discovery of such large \mathcal{B})



Branching fraction of charmed baryon decays







- The weak decays of charmed baryons provide an excellent platform for understanding QCD with transitions involving the charm quark. The decay amplitudes consist of factorizable and non-factorizable contributions.
- First or most precise \mathcal{B} results for charmed baryon decays.

- CF decays:

- $\mathcal{B}(\Lambda_c^+ \rightarrow p K_S^0 \eta) = (4.35 \pm 0.10 \pm 0.20 \pm 0.22) \times 10^{-3}$
- $\mathcal{B}(\Lambda_c^+ \rightarrow \Sigma^+ \eta) = (3.14 \pm 0.35 \pm 0.17 \pm 0.25) \times 10^{-3}$
- $\mathcal{B}(\Lambda_c^+ \rightarrow \Sigma^+ \eta') = (4.16 \pm 0.75 \pm 0.25 \pm 0.33) \times 10^{-3}$

- SCS and DCS decays:

- $\mathcal{B}(\Lambda_c^+ \rightarrow \Lambda K^+) = (6.57 \pm 0.17 \pm 0.11 \pm 0.35) \times 10^{-4}$
- $\mathcal{B}(\Lambda_c^+ \rightarrow \Lambda K^+) = (3.58 \pm 0.19 \pm 0.06 \pm 0.19) \times 10^{-4}$
- $\mathcal{B}(\Lambda_c^+ \rightarrow p K_S^0 K_S^0) = (2.35 \pm 0.12 \pm 0.07 \pm 0.12) \times 10^{-4}$
- $\frac{\mathcal{B}(\Lambda_c^0 \rightarrow \Xi^- \pi^+)}{\mathcal{B}(\Lambda_c^0 \rightarrow \Omega^- \pi^+)} = 0.253 \pm 0.052 \pm 0.030$
- $\frac{\mathcal{B}(\Lambda_c^0 \rightarrow \Xi^- K^+)}{\mathcal{B}(\Lambda_c^0 \rightarrow \Omega^- \pi^+)} < 0.070$



Decay asymmetry parameter α of charmed baryon decays

- The **decay asymmetry parameter α** was introduced by Lee and Yang to study the parity-violating and parity-conserving amplitudes in weak hyperon decays.
- In $1/2^+ \rightarrow 1/2^+ + 0^-$, $\alpha \equiv 2 \cdot \text{Re}(S^*P) / (|S|^2 + |P|^2)$, where S and P denote the parity-violating S -wave and parity-conserving P -wave amplitudes.
- The charmed baryons are produced with negligible polarization averagely at Belle (II), different with BESIII/LHCb.
- For $\Lambda_c^+ \rightarrow \Lambda h^+, \Sigma^+ h^0$ decays, the differential decay rate depends on α :


$$\frac{dN(\Lambda_c^+ \rightarrow \Lambda h^+)}{d \cos \theta_\Lambda} \propto 1 + \alpha_{\Lambda_c^+} \alpha_- \cos \theta_\Lambda$$

where α_- is hyperon decay asymmetry parameter.

- For $\Lambda_c^+ \rightarrow \Sigma^0 h^+$ decays, considering $\alpha(\Sigma^0 \rightarrow \gamma \Lambda)$ is zero due to parity conservation for an electromagnetic decay, the differential decay rate


$$\frac{dN(\Lambda_c^+ \rightarrow \Sigma^0 h^+)}{d \cos \theta_{\Sigma^0} d \cos \theta_\Lambda} \propto 1 - \alpha_{\Lambda_c^+} \alpha_- \cos \theta_{\Sigma^0} \cos \theta_\Lambda$$

- By studying the hyperon helicity angle, we can extract α of charmed baryon decays.



Decay asymmetry parameter α of charmed baryon decays

- The efficiency-corrected yields are used to extract α .



- No approaches based on various theories could successfully predict all these experimental α values.
 ⇒ needs a joint effort from theory and experiment in future.

Decay	α at Belle	W.A. or BESIII
$\Lambda_c^+ \rightarrow p K_S^0$	-	0.18 ± 0.45^a
$\Lambda_c^+ \rightarrow \Lambda K^+$	-0.585 ± 0.052^b	-
$\Lambda_c^+ \rightarrow \Sigma^0 K^+$	-0.54 ± 0.20^b	-
$\Lambda_c^+ \rightarrow \Lambda \pi^+$	-0.755 ± 0.006^b	-0.84 ± 0.09
$\Lambda_c^+ \rightarrow \Sigma^0 \pi^+$	-0.463 ± 0.018^b	-0.73 ± 0.18^c
$\Lambda_c^+ \rightarrow \Sigma^+ \pi^0$	-0.480 ± 0.028^d	-0.55 ± 0.11
$\Lambda_c^+ \rightarrow \Sigma^+ \eta$	-0.990 ± 0.058^d	-
$\Lambda_c^+ \rightarrow \Sigma^+ \eta'$	-0.460 ± 0.067^d	-
$\Lambda_c^+ \rightarrow \Xi^0 K^+$	-	0.01 ± 0.16^e
$\Lambda_c^+ \rightarrow \Lambda \rho^+$	-	-0.76 ± 0.07^f
$\Lambda_c^+ \rightarrow \Sigma^{*+} \pi^0$	-	-0.92 ± 0.09^f
$\Lambda_c^+ \rightarrow \Sigma^{*0} \pi^+$	-	-0.79 ± 0.11^f
<hr/>		
$\Xi_c^0 \rightarrow \Xi^- \pi^+$	-0.64 ± 0.05^g	$-0.56 \pm 0.39^{+0.10}_{-0.09}^h$
$\Xi_c^0 \rightarrow \Lambda \bar{K}^{*0}$	$+0.15 \pm 0.22^i$	-
$\Xi_c^0 \rightarrow \Sigma^+ K^{*-}$	-0.52 ± 0.30^h	-

^aBESIII, PRD 100, 072004 (2019)

^bBelle, Sci. Bull. 68, 583 (2023)

^cBESIII, PRD 100, 072004 (2019)

^dBelle, PRD 107, 032003 (2023)

^eBESIII, arXiv:2309.02774

^fBESIII, JHEP 12, 033 (2022)

^gBelle, PRL 127, 121803 (2021)

^hCLEO, PRD 63, 111102 (2001)

ⁱBelle, JHEP 06, 160 (2021)




Outline

- 1 Charm sample at Belle and Belle II
- 2 Charm lifetime measurements
- 3 Branching fraction and decay asymmetry parameter
- 4 Search for rare or forbidden decay**
- 5 Charm CP violation searches
- 6 Summary

Search for $\Xi_c^0 \rightarrow \Xi^0 \ell^+ \ell^-$ at Belle

(Belle) arXiv:2312.02580

- In the Standard Model (SM), the weak-current interaction has an identical coupling to all lepton generations, which allows Lepton Flavor Universality (LFU) to be tested in the semileptonic decays of the hadrons.
- The $\Xi_c^0 \rightarrow \Xi^0 \ell^+ \ell^-$ are related to the W -exchange contribution in $\Lambda_c^+ \rightarrow p \ell^+ \ell^-$.
- Measurement of both $\ell = e$ channel and μ decay rates would allow an LFU test to be performed
- Result: $\frac{\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- e^+ e^-)}{\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)}$ $< 6.7 \times 10^{-3}$ and $\frac{\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \mu^+ \mu^-)}{\mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+)}$ $< 4.3 \times 10^{-3}$
- A more precise analysis based on larger data samples collected by Belle II is expected in the future.




Search for $D \rightarrow p\ell$ at Belle

(Belle) arXiv:2310.07412

- Baryon number violation (BNV) is one of the crucial conditions to create the matter-antimatter asymmetry as observed in the universe.
- Several grand unified theories, supersymmetry and other SM extensions propose BNV processes of nucleons.
- $D \rightarrow p\ell$: baryon (B) and lepton (L) numbers violated but their difference conserved ($\Delta(B-L) = 0$)
- pre-Belle stringent limits: $\mathcal{B}(D^0 \rightarrow \bar{p}e^+) < 1.2 \times 10^{-6}$ and $\mathcal{B}(D^0 \rightarrow pe^-) < 2.2 \times 10^{-6}$ from BESIII.
- Recently, Belle set their upper limits in the range $(5-8) \times 10^{-7}$ at a 90% C.L., depending on the decay mode:

TABLE I. Reconstruction efficiency (ϵ), signal yield (N_S), signal significance (S), upper limit on the signal yield (N_{pl}^{UL}), and branching fraction (\mathcal{B}) at 90% confidence level for each decay mode.

Decay mode	ϵ (%)	N_S	S (σ)	N_{pl}^{UL}	$\mathcal{B} \times 10^{-7}$
$D^0 \rightarrow pe^-$	10.2	-6.4 ± 8.5	—	17.5	< 5.5
$\bar{D}^0 \rightarrow pe^-$	10.2	-18.4 ± 23.0	—	22.0	< 6.9
$D^0 \rightarrow \bar{p}e^+$	9.7	-4.7 ± 23.0	—	22.0	< 7.2
$\bar{D}^0 \rightarrow \bar{p}e^+$	9.6	7.1 ± 9.0	0.6	23.0	< 7.6
$D^0 \rightarrow p\mu^-$	10.7	11.0 ± 23.0	0.9	17.1	< 5.1
$\bar{D}^0 \rightarrow p\mu^-$	10.7	-10.8 ± 27.0	—	21.8	< 6.5
$D^0 \rightarrow \bar{p}\mu^+$	10.5	-4.5 ± 14.0	—	21.1	< 6.3
$\bar{D}^0 \rightarrow \bar{p}\mu^+$	10.4	16.7 ± 8.8	1.6	21.4	< 6.5



Outline


- 1 Charm sample at Belle and Belle II
- 2 Charm lifetime measurements
- 3 Branching fraction and decay asymmetry parameter
- 4 Search for rare or forbidden decay
- 5 Charm CP violation searches**
- 6 Summary



Why CP Violation and Charm CPV Special?

- Standard Model provides a solo CP violation (CPV) source in quark sector: a complex phase in CKM matrix. But, such CPV source is not large enough to explain the observed matter-antimatter asymmetry of the universe.
 - ⇒ **search for new CPV sources beyond SM**, as a lasting hot topic.
 - Sakharov in 1967: CPV is one of the three conditions necessary to explain the matter-antimatter asymmetry of the universe.
 - In 1964, CPV in K meson decays was observed by Cronin, Fitch, *et al.* (Nobel 1980)
 - In 2001, BABAR and Belle observed a large CP asymmetry in B meson decays, providing strong experimental evidence for the theoretical predictions from Kobayashi and Maskawa (Nobel 2008).
- Charm CPV effect is very small ($\mathcal{O}(10^{-3})$ or smaller ^{*ab*}). New Physics may enhance it ^{*cd*}.
- Study of charm CPV may help to understand the SM, and is a sensitive probe to search for New Physics.
- In 2019, CP violation in D^0 decays was found at LHCb: $\Delta A_{CP}(D^0 \rightarrow K^+K^-, \pi^+\pi^-) = (-15.4 \pm 2.9) \times 10^{-4}$ (5.3σ).
 - ⇒ to understand this CPV, **we need to study more channels and improve the precision on the existing measurements.**
- CPV has been observed in all the open-flavored meson sector, but **not yet established in the baryon sector.**
 - ⇒ **discovering the CPV in charmed baryon is one of major targets of charm physics.**

Citations per year



^aH.-n. Li, C.-D. Lu, and F.-S. Yu, [PRD 86, 036012 \(2012\)](#)

^bH.-Y. Cheng and C.-W. Chiang, [PRD 104, 073003 \(2021\)](#)

^cA. Dery and Y. Nir, [JHEP 12, 104 \(2019\)](#)

^dM. Saur and F.-S. Yu, [Sci. Bull. 65, 1428 \(2020\)](#)

CPV in charm four-body decays


- Sensitivity of CPV relies on various physical decays, motivating the exploration of CPV in diverse charm decays.
- The D four-body decay with large β and various intermediate processes provides different sample for CPV searches.
- CPV in D four-body decay was searched with triple-product asymmetries by the FOCUS, BABAR, LHCb and Belle experiments.
- Triple-product $C_T = (\vec{p}_1 \times \vec{p}_2) \cdot \vec{p}_3$ is calculated in the mother's rest frame, satisfying $CP(C_T) = -C(C_T) = -\bar{C}_T$.

- The T-odd asymmetries, taking for D^\pm decays for example, are defined as

$$A_T(D^+) = \frac{\Gamma_+(C_T > 0) - \Gamma_+(C_T < 0)}{\Gamma_+(C_T > 0) + \Gamma_+(C_T < 0)} \quad \bar{A}_T(D^-) = \frac{\Gamma_-(-\bar{C}_T > 0) - \Gamma_-(-\bar{C}_T < 0)}{\Gamma_-(-\bar{C}_T > 0) + \Gamma_-(-\bar{C}_T < 0)}$$

- T-odd CP-asymmetry $a_{CP}^{T\text{-odd}} = \frac{1}{2}(A_T - \bar{A}_T)$, may be an observable complementary to the direct CPV (A_{CP}^{dir})



With some conditions, $a_{CP}^{T\text{-odd}} \propto \sin\phi \cos\delta$ has largest value when $\delta = 0$ ($A_{CP}^{\text{dir}} \neq 0$ needs $\delta \neq 0$)



CPV in charm four-body decays

(Belle) PRD 107, 052001 (2023); PRD 108, L111102 (2023); arXiv:2305.12806

- Belle recently searched for CPV with T -odd correlations in decays of $D^0 \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$, $D_{(s)}^+ \rightarrow K_S^0 h^+ \pi^+ \pi^-$ and $D_{(s)}^+ \rightarrow Kh \pi^+ \pi^0$.
- These $a_{CP}^{T\text{-odd}}$ results mostly are first or most precise measurement.



- $\sigma(a_{CP}^{T\text{-odd}})$ of all D mesons: reached $\mathcal{O}(0.1\%)$ level.
- Belle II/LHCb may improve the precision utilizing increased samples, and apply this method to charmed baryons.

direct CPV in $\Lambda_c^+ \rightarrow \Lambda K^+, \Sigma^0 K^+$

(Belle) Science Bulletin 68 (2023) 583


- The raw asymmetry of $\Lambda_c^+ \rightarrow \Lambda K^+$ includes several asymmetry sources:

$$A_{\text{raw}}(\Lambda_c^+ \rightarrow \Lambda K^+) \approx A_{CP}^{\Lambda_c^+ \rightarrow \Lambda K^+} + A_{CP}^{\Lambda \rightarrow p\pi^-} + A_\epsilon^\Lambda + A_\epsilon^{K^+} + A_{FB}^{\Lambda_c^+}$$

- $A_{CP}^{\Lambda_c^+ \rightarrow \Lambda K^+}$ ($A_{CP}^{\Lambda \rightarrow p\pi^-}$): direct CP asymmetry associated with Λ_c^+ (Λ) decay,
- A_ϵ^Λ ($A_\epsilon^{K^+}$): detection asymmetry arising from efficiencies between Λ (K^+) and $\bar{\Lambda}$ (K^-). The $A_\epsilon^{K^+}$ is removed by weighting $w_{\Lambda_c^+, \bar{\Lambda}_c^-} = 1 \mp A_\epsilon^{K^+} [\cos\theta, p_T]$
- $A_{FB}^{\Lambda_c^+}$ arises from the forward-backward asymmetry (FBA) of Λ_c^+ production due to γ - Z^0 interference and higher-order QED effects in $e^+e^- \rightarrow c\bar{c}$ collisions. The FBA is an odd function in $\cos\theta^*$, where θ^* is the Λ_c^+ production polar angle in the e^+e^- center-of-mass frame, but due to asymmetric acceptance, small residual asymmetry remains after integrating over $\cos\theta^*$.
- using CF mode $\Lambda_c^+ \rightarrow \Lambda\pi^+$ to remove the common asymmetry sources.

- Result: $\Delta A_{\text{raw}} = A_{\text{raw}}^{\text{corr}}(\Lambda_c^+ \rightarrow \Lambda K^+) - A_{\text{raw}}^{\text{corr}}(\Lambda_c^+ \rightarrow \Lambda\pi^+) = A_{CP}^{\text{dir}}(\Lambda_c^+ \rightarrow \Lambda K^+) - A_{CP}^{\text{dir}}(\Lambda_c^+ \rightarrow \Lambda\pi^+) = A_{CP}^{\text{dir}}(\Lambda_c^+ \rightarrow \Lambda K^+)$

The reference mode and signal mode have nearly same Λ kinematic distributions, including the Λ decay length, the polar angle and the momentum of the proton and pion in the laboratory reference frame.



- $A_{CP}^{\text{dir}}(\Lambda_c^+ \rightarrow \Lambda K^+) = (+2.1 \pm 2.6 \pm 0.1)\%$
- $A_{CP}^{\text{dir}}(\Lambda_c^+ \rightarrow \Sigma^0 K^+) = (+2.5 \pm 5.4 \pm 0.4)\%$


First A_{CP}^{dir} for SCS two-body decays of charmed baryons.

baryonic α -induced CPV in $\Lambda_c^+ \rightarrow \Lambda K^+, \Sigma^0 K^+$

(Belle) Science Bulletin 68 (2023) 583

(SCS) $\Lambda_c^+ \rightarrow \Lambda K^+, \Sigma^0 K^+$


- Measure $\alpha/\bar{\alpha}$ for the separate $\Lambda_c^+/\bar{\Lambda}_c^-$ samples.
- Calculate $A_{CP}^\alpha \equiv (\alpha_{\Lambda_c^+} + \alpha_{\bar{\Lambda}_c^-})/(\alpha_{\Lambda_c^+} - \alpha_{\bar{\Lambda}_c^-})$.



- Result: $A_{CP}^\alpha(\Lambda_c^+ \rightarrow \Lambda K^+) = -0.023 \pm 0.086 \pm 0.071$
 $A_{CP}^\alpha(\Lambda_c^+ \rightarrow \Sigma^0 K^+) = +0.08 \pm 0.35 \pm 0.14$
First A_{CP}^α results for charmed baryon SCS decays.
- No evidence of CPV is found.

(CF) $\Lambda_c^+ \rightarrow \Lambda \pi^+, \Sigma^0 \pi^+$

- Probe Λ -hyperon CPV in charmed baryon CF decays, inspired by [arXiv:2208.01589](https://arxiv.org/abs/2208.01589).
- Under a reasonable assumption $\alpha_{\Lambda_c^+} = -\alpha_{\bar{\Lambda}_c^-}$ in CF decays, we have $A_{CP}^\alpha(\Lambda \rightarrow p\pi^-) = A_{CP}^\alpha(\text{total}) \equiv \frac{\alpha_{\Lambda_c^+} + \alpha_{\bar{\Lambda}_c^-}}{\alpha_{\Lambda_c^+} - \alpha_{\bar{\Lambda}_c^-}}$.



- Result: $A_{CP}^\alpha(\Lambda \rightarrow p\pi^-) = +0.013 \pm 0.007 \pm 0.011$
The first result of hyperon CPV in charm CF decays


Outline

- 1 Charm sample at Belle and Belle II
- 2 Charm lifetime measurements
- 3 Branching fraction and decay asymmetry parameter
- 4 Search for rare or forbidden decay
- 5 Charm CP violation searches
- 6 Summary**



Summary

- **Belle are lasting to produce the fruitful charm results** although its data taking finished 13 years ago. My talk selected some recent results on B and α measurements and CPV searches for charm decays.
- **Belle II has joined the game.** To date, Belle II has collected 424 fb^{-1} of dataset.
- Utilizing the early dataset, we obtain the world best $\tau(D^{0,+})$, $\tau(D_s^+)$, and $\tau(\Lambda_c^+)$ (first Belle II precision measurements) and confirmation of LHCb $\tau(\Omega_c^0)$ result are presented.
- Stay tuned for more charm results utilizing 1.4 ab^{-1} of dataset at Belle and Belle II in 2024, and more luminosity (goal: 50 ab^{-1}) in the future at Belle II.



Some homework of charm physics in my personal superficial opinion

- first observation of charm CPV in singly decay channel and more channels of D mesons

$$\Delta A_{CP}(D^0 \rightarrow K^+K^-, \pi^+\pi^-) (> 5\sigma) \text{ and } A_{CP}^{\text{dir}}(D^0 \rightarrow \pi^+\pi^-) (3.8\sigma)$$

CP asymmetry in many SCS decay channels have been studied but with statistics limited.

- first evidence of indirect CPV in D^0 decays [Long term]

still no signs for non-zero result in $|q/p| - 1$ and $\arg(q/p)$.

- first evidence of CPV in charmed baryon sector [Long term]

$$\text{currently only three studies } \Lambda_c^{\pm} \rightarrow ph^+h^-, \Lambda_c^{\pm} \rightarrow (\Lambda, \Sigma^0)K^{\pm}, \Xi_c^{\pm} \rightarrow pK^-\pi^{\pm}$$

- first observation of Ξ_{cc}^+ and Ω_{cc}^+ and their hadronic decays

- first observation of radiative decays of charmed baryons

- precise/first absolute \mathcal{B} of the decays of charmed baryons (Ξ_c and Ω_c)

- more precise \mathcal{B} results of charmed baryon SL decays


e.g $\mathcal{B}(\Xi_c \rightarrow \Xi \ell \nu)$ and $\mathcal{B}(\Omega_c \rightarrow \Omega \ell \nu)$ results are not understood or to be improved precisely.

- \mathcal{B} (and α) measurements for more charm decays or with improved precision

- amplitude analyses of charmed baryon decays with current/increased available datasets

- more sensitive searches for rare or forbidden charm decays [Long term]

-




"The road ahead is long and endless; yet high and low we'll search with our will unbending."

Thank you for your attentions.



谢谢!

Dr. Longke LI (李龙科)
Department of Physics,
University of Cincinnati (UC)

 lilongke_ustc

 lilk@ucmail.uc.edu




Why measure charm lifetimes?

Lenz, IJMP A30 (2015)
 Lenz et al., JHEP 12 (2020) 199
 King, Lenz et al., JHEP 08 (2022) 241
 Gratx et al., JHEP 07 (2022) 058

Theory:

- **qualitatively understood in terms of simple diagrams,**
 e.g., $c \rightarrow s e^+ \nu$ partial width gives $G_F^2 m_c^5 |V_{cs}|^2 / (192\pi^3)$ dependence. Long D^+ lifetime can be understood as arising from destructive interference between spectator and color-suppressed amplitudes. But this doesn't include QCD...
- **to include QCD: calculate using the Heavy Quark Expansion**



$$\Gamma(D) = \frac{1}{2m_D} \sum_X \int_{PS} (2\pi)^4 \delta^4(p_D - p_X) |\langle X(p_X) | \mathcal{H}_{\text{eff}} | D(p_D) \rangle|^2,$$

ΣX is sum over final states

$$\rightarrow \frac{1}{2m_D} \text{Im} \langle D | \mathcal{T} | D \rangle \quad \text{where} \quad \mathcal{T} = i \int d^4x T \{ \mathcal{H}_{\text{eff}}(x), \mathcal{H}_{\text{eff}}(0) \}$$


via optical theorem

$$\rightarrow \Gamma_3 + \Gamma_5 \frac{\langle \mathcal{O}_5 \rangle}{m_c^2} + \Gamma_6 \frac{\langle \mathcal{O}_6 \rangle}{m_c^3} + \dots + 16\pi^2 \left(\tilde{\Gamma}_6 \frac{\langle \tilde{\mathcal{O}}_6 \rangle}{m_c^3} + \tilde{\Gamma}_7 \frac{\langle \tilde{\mathcal{O}}_7 \rangle}{m_c^4} + \dots \right)$$

via Heavy Quark Expansion

Wilson coefficients Γ_i are expanded in powers of α_s and calculated perturbatively





\Rightarrow comparing lifetime calculations with measurements tests/improves our understanding of QCD




from KEKB to SuperKEKB

(For more info, see [Xiaolong's talk](#))

- ▶ As 1st and 2nd generation B-factories, KEKB and SuperKEKB have many similarities, and more differences:
 - Damping ring added to have low emittance positrons / use 'Nano-beam' scheme by squeezing the beta function at the IP.
 - beam energy: admit lower asymmetry to mitigate Touschek effects / beam current: $\times 2$ to contribute to higher luminosity.
 - SuperKEKB achieved the luminosity record of $4.7 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$.



Detector: Belle II Vs. Belle



SVD: 4 lysrs \rightarrow VXD=(PXD 2 lysrs + SVD 4 lysrs)
 CDC: small cell, long lever arm
 ACC+TOF \rightarrow TOP+ARICH
 ECL: waveform sampling
 KLM: RPC \rightarrow Scintillator + SiPM
 (endcaps, barrel inner 2 lysrs)

