





# Belle II and anomalies: what's new, what's next?

**Vitalii Lisovskyi** (Aix Marseille Univ, CNRS/IN2P3, CPPM)  
on behalf of the Belle II collaboration

**Siegen, 11 April 2024**


# The journey so far

- Belle II: so far, mostly running on/near  $\Upsilon(4S)$  resonance, with a short scan above  $\Upsilon(4S)$
- Can include the Belle data:  $\Upsilon(1 - 5S)$



# The journey so far


- World record peak luminosity in summer 2022:  $4.7 \times 10^{34}/\text{cm}^2/\text{s}$
- Progress limited by sudden beam losses and other accelerator issues
- LS1: machine improvements & complete the vertex detector
- **The data taking has resumed recently**
- Main objectives for 2024:
  - Reach and maintain the peak luminosity of  $10^{35}/\text{cm}^2/\text{s}$
  - Cross the  $1\text{ab}^{-1}$  milestone
  - Prove the effectiveness of the work done in LS1



Not just more data: improvement in the analysis tools




# Improving the tagging: beauty

- **Conventional category-based tagger** (by charge of the lepton/kaon from the other B): effective tagging efficiency  $\sim 30\%$
- Flavour tagging using machine-learning techniques with the full event information (PID, tracking, kinematics) for the "rest of the event"
  - New **GNN-based tagger GFlaT** [[2402.17260](#)] with 37% efficiency
  - Works best when leptons and/or kaons present in the ROE, less well for pion-only events
- Prospects for improvement: requires better understanding of simulation




# Improving the tagging: charm

- **Conventional method:** tag the charge of the pion from  $D^{*\pm} \rightarrow D^0\pi^\pm$ , or lepton from  $B^- \rightarrow D^0\ell^-\nu$ 
  - Loss of statistics due to low production rates, soft pion efficiency etc: ~25% effective efficiency
- New **inclusive tagging** in  $e^+e^- \rightarrow c\bar{c}$  with BDT algorithm [[Phys. Rev. D 107, 112010 \(2023\)](#)]
  - Uses OS and SS information
  - Effective tagging eff. ~48%
  - **Doubles the effective sample size** for CPV and charm mixing studies
  - Useful to **suppress backgrounds** in untagged analyses




# Missing-energy estimation

- **Conventional approach:** full reconstruction of the "other B" in a number of specific decay modes
  - Full event interpretation
  - Hadronic or semileptonic tag
- **Inclusive tagger:** reconstruct signal first, use the "rest of the event" for tagging
- Disagreement between data and simulated performance needs to be calibrated
- Constant improvements:
  - New tag decay modes added
  - BF and resonant structures of known decays re-measured to improve the simulation
  - Alternative ML tools explored (graph neural networks...)



# Further ahead

- **The eventual target is to collect 50 ab<sup>-1</sup>, but:**
- Going beyond  $2 \times 10^{35}/\text{cm}^2/\text{s}$  requires a redesign of the interaction region and the vertex detector: beam background
- Envisaged LS2 in 2027-2028, no precise planning yet
- May profit from this shutdown for other detector improvements
- The priority is to run at/near  $\Upsilon(4S)$ ; special datasets at different energies might be collected in the future




# *A biased collection of physics topics*

Focus mostly on the topics **not** covered by Florian, Markus and Caspar

For physics prospects, a recent reference is the 2022 Snowmass report: [2207.06307](#)



# Belle II versus anomalies (broadly defined)



# Tau properties

$$\mathcal{B}(\tau \rightarrow e\nu\bar{\nu}) \sim \mathcal{B}(\mu \rightarrow e\nu\bar{\nu}) \frac{m_\tau^5}{m_\mu^5} \quad 10$$




- Abundant  $e^+e^- \rightarrow \tau^+\tau^-$  production

- **Tau mass:** pseudomass method with  $e^+e^- \rightarrow \tau^+\tau^-$ ,  $\tau \rightarrow 3\pi\nu$   
 $1777.09 \pm 0.08 \pm 0.11 \text{ MeV}/c^2$   
 most precise to date  
 [Phys. Rev. D 108, 032006]

- Largest syst: beam energy scale, momentum scale
  - Affected by the knowledge of  $\Upsilon(4S)$  lineshape and B mass!

- Next step: **tau lifetime** (Belle result is world best)

- Belle II will reduce both stat. and syst. uncertainties significantly, down to  $0.2 \times 10^{-15} \text{ s}$



- **Lepton universality** in tau decays:

- $\frac{\mathcal{B}(\tau \rightarrow \mu\nu\bar{\nu})}{\mathcal{B}(\tau \rightarrow e\nu\bar{\nu})}$  (mu/e universality) and  $\frac{\Gamma(\tau \rightarrow \pi\nu)}{\Gamma(\pi \rightarrow \mu\nu)}$  (mu/tau universality)

- Sensitivity eventually limited by the control of PID performance, but can improve world average by a factor of ~few.

- **Michel parameters** (Lorentz structure of the  $\tau \rightarrow \mu\nu\nu$  decay):

kink reconstruction at Belle allows to measure with ~100% uncertainty, precision down to few % expected with new algorithms & enlarged drift chamber at Belle II



- Input to the **Cabibbo angle anomaly**:  $|V_{us}|$  from  $\frac{\mathcal{B}(\tau \rightarrow K\nu)}{\mathcal{B}(\tau \rightarrow \pi\nu)}$

- Projected reach down to ~1% sensitivity, depending on PID performance

- CPV measurements in tau decays...

# LFV searches in $\tau$ decays

- The most stringent upper limits on **lepton-flavour-violating tau decays** come from the B factories
- Recent Belle II  $\tau \rightarrow 3\mu$  search:
  - Tag  $e^+e^- \rightarrow \tau^+\tau^-$  with 1-3 tracks on the tag side
  - Look for events with  $E_{sig} - E_{beam} = 0$  near the  $\tau$  mass
  - Efficiency 3x better than Belle!
  - UL:  $\mathcal{B} < 1.9 \times 10^{-8}$ @90%CL
- Belle II projected reach:  $(5 - 10) \times 10^{-10}$  for  $50 \text{ ab}^{-1}$  for most LFV channels
  - except for  $\tau \rightarrow \ell\gamma$  modes (irreducible bkg due to  $\tau \rightarrow \ell\nu\bar{\nu} + \gamma_{ISR}$ )
    - This is where beam polarisation may help (about this later)
- Analyses of many other final states in progress
  - Also, searches for new bosons in  $\tau \rightarrow \ell a$ , HNL in  $\tau \rightarrow \pi N$ ...




- The recent  $B^+ \rightarrow K^+ \nu \bar{\nu}$  analysis presented by Caspar. Expect observation with more data!
  - Clarifying the background properties is important
  - Future prospects:  $B \rightarrow K^* \nu \bar{\nu}$ ,  $B \rightarrow K_S^0 \nu \bar{\nu}$ , inclusive  $B \rightarrow X_S \nu \bar{\nu}$ ,  $B \rightarrow \pi(\rho) \nu \bar{\nu}$ ...
    - Spin-offs: BSM searches in  $B \rightarrow K^{(*)} + \text{invisible}$ , charm decays e.g.  $\Lambda_c^+ \rightarrow p \nu \bar{\nu}$  or  $D \rightarrow \pi \nu \bar{\nu}$  (GIM-suppressed)
- **Experimental techniques (missing energy) can be applied to  $B \rightarrow K^* \tau^+ \tau^-$ ,  $B \rightarrow \rho \tau^+ \tau^-$  searches**
  - Expected sensitivity down to  $5 \times 10^{-4}$  BF, still far away from the SM rate
  - As well as LFV  $b \rightarrow s \tau^\pm \ell^\mp$  searches, with sensitivity down to few  $\times 10^{-6}$
- Let me reiterate the importance of understanding & improving the tagging performance

- Reach in  $b \rightarrow s(d)e^+e^-$  and  $b \rightarrow s(d)\mu^+\mu^-$  statistically limited compared to LHCb, but similar performance in muons and electrons helps for LFU tests.
- Very competitive in final states with neutrals e.g.  $B^0 \rightarrow \pi^0 e^+ e^-$ , see [recent Belle result](#)

	$N_{\text{sig}}$	$\mathcal{B}^{\text{UL}} (10^{-8})$
$B^0 \rightarrow \eta \ell^+ \ell^-$	$0.5^{+1.0}_{-0.8}$	$< 4.8$
$B^0 \rightarrow \eta e^+ e^-$	$0.0^{+1.4}_{-1.0}$	$< 10.5$
$B^0 \rightarrow \eta \mu^+ \mu^-$	$0.8^{+1.5}_{-1.1}$	$< 9.4$
$B^+ \rightarrow \pi^+ e^+ e^-$	$0.1^{+2.5}_{-1.6}$	$< 5.4$
$B^0 \rightarrow \pi^0 \ell^+ \ell^-$	$-1.8^{+1.6}_{-1.1}$	$< 3.8$
$B^0 \rightarrow \pi^0 e^+ e^-$	$-2.9^{+1.8}_{-1.4}$	$< 7.9$
$B^0 \rightarrow \pi^0 \mu^+ \mu^-$	$-0.5^{+3.6}_{-2.7}$	$< 5.9$


- Belle II is crucial to provide the measurements of **absolute branching fractions**, e.g. for normalisation modes used by LHCb such as  $B^+ \rightarrow K^+ J/\psi$

- Recent Belle+Belle II search for  $B^0 \rightarrow \gamma\gamma$  (see Moriond talk) - a very suppressed  $b \rightarrow d$  transition
- $UL < 6.4 \times 10^{-8}$  @90% CL, only factor  $\sim 5$  above the SM prediction
- Very interesting measurement with  $50ab^{-1}$ !
- $D^0 \rightarrow \gamma\gamma$  search in prospects: sensitivity down to  $\sim 10^{-7}$  (factor  $\sim 10$  above the SM rate)
- A less suppressed  $B_s^0 \rightarrow \gamma\gamma$  can be searched if  $B_s^0$  data collected



# Radiative (charm) decays


- Many measurements of  $b \rightarrow s\gamma$ : e.g. **photon polarisation** in  $B \rightarrow K\pi\pi\gamma$  down to  $\sim 1\%$  at  $50 \text{ ab}^{-1}$
- In the **charm sector**, the penguin  $c \rightarrow u\gamma$  is very suppressed
  - The  $4\pi$  geometry of Belle (II) helps with rejecting  $c \rightarrow u\pi^0$  backgrounds
  - W exchange  $cd \rightarrow us\gamma$  (long-distance) is expected to have a larger rate
    - Interest to measure photon polarisation
- Belle did the first search for **radiative charm baryon decays**  $\Lambda_c^+ \rightarrow \Sigma^+\gamma$  and  $\Xi_c^0 \rightarrow \Xi^0\gamma$  [[Phys. Rev. D 107, 032001 \(2023\)](#)]
- BF limits at the  $2 \times 10^{-4}$  level, hope for observation with Belle II data?
  - Theory predictions in  $\text{few} \times 10^{-5}$  range







# A few words on $b \rightarrow c\ell\nu$ and $b \rightarrow u\ell\nu$

- Closing the gap between inclusive and exclusive decays to corner the  $|V_{cb}|$




- $|V_{ub}|$  inclusive down to ~3-5% precision with 50ab<sup>-1</sup> (theory-dominated), exclusive more precise
- Precision on  $R_{D^{(*)}}$  down to few %
- High hope to observe  $B^+ \rightarrow \mu^+ \nu$  and improve significantly  $B^+ \rightarrow \tau^+ \nu$  measurement: both down to ~5% relative unc. with 50ab<sup>-1</sup>
  - Benefit from inclusive tagging developed for  $B^+ \rightarrow K^+ \nu \nu$
  - Don't forget about  $B^+ \rightarrow \mu^+ \nu \gamma$  (see [here](#))



# CP violation in B decays







- **We enter the era of precision testing of the CKM unitarity**
- The unique feature of Belle II is the entangled B production and, therefore, high flavour-tagging efficiency
  - NB: search for non-perfect entanglement is an interesting QM test!
- World-best sensitivity achievable in final states with  $\pi^0$ ,  $K_L^0$  or  $K_S^0$
- At 50  $\text{ab}^{-1}$ , expected precision of
  - $<1\%$  on  $\sin 2\phi_1^{(eff)} \equiv \sin 2\beta^{(eff)}$  in tree-dominated  $(c\bar{c})K^0$  or  $\sim 1.5\%$  in loop-dominated  $\eta'K^0$ 
    - See the recent result in  $B^0 \rightarrow J/\psi K_S^0$  with early Belle II data (3x worse than LHCb Run1+2)
  - $\sim 2^\circ$  on  $\phi_2 \equiv \alpha$  in  $B \rightarrow \rho\rho$
  - $\sim 2^\circ$  on  $\phi_2 \equiv \gamma$
- Narrowing down on the isospin sum rule in  $B \rightarrow K\pi$  decays ("K $\pi$  puzzle"), where  $A_{CP}(B \rightarrow K_S^0\pi^0)$  will be driven by Belle II (down to few % at 50  $\text{ab}^{-1}$ )
  - Recent Belle II result compatible with the SM:  $I_{K\pi} = -0.03 \pm 0.13 \pm 0.04$
- CPV studies in charmless B decays dominated by Belle II



- Belle II uniquely positioned to probe CPV in final states with neutrals
- $D^0 \rightarrow \pi^0\pi^0$  and  $D^+ \rightarrow \pi^+\pi^0$  are well motivated
  - Sensitivity down to 0.07% ( $50 \text{ ab}^{-1}$ ) for  $D^0 \rightarrow \pi^0\pi^0$  with the conventional  $D^*$  tag
    - but we have a much better tagger now!
- Isospin **sum rule** by comparing CPV in  $D^0 \rightarrow \pi^+\pi^-$ ,  $D^0 \rightarrow \pi^0\pi^0$  and  $D^+ \rightarrow \pi^+\pi^0$  decays: probe whether CPV is SM or beyond

# Lifetimes


- Decay-time resolution 2x better than Belle
- Fewer sculpting effects compared to the LHCb trigger



World best for all except  $\Omega_c^0$  and still statistically limited!



Much more precise than theory

Systematics: detector alignment & resolution, backgrounds




# Inclusive charm baryons

- Two ways to obtain inclusive charm baryon datasets:
  - baryon-number,  $s$  and  $c$  conservation in  $e^+e^-$  collisions:
 
$$e^+e^- \rightarrow D^{(*)-}\bar{p}\pi^+\Lambda_c^+$$
    - notable example: [[Phys.Rev.Lett. 113 \(2014\) 4, 042002](#)]
  - B-meson decays,  $\bar{B}^0 \rightarrow \Xi_c^+\bar{\Lambda}_c^-$  with one baryon treated as recoi
    - notable example: [[Phys.Rev.D 100 \(2019\) 3, 031101](#)]
    - low statistics
- Useful to measure absolute BF, but in particular **decays with missing energy (semileptonic)**
- More results expected with these methods
  - Absolute BFs of  $\Xi_c/\Omega_c$  imprecise or unknown (more data & better tagging helps!)
  - Note: BES III catching up by running on baryon-pair thresholds



# Dipole moments

- **Muon g-2 inputs:** measure cross-section inputs to the HVP calculation
  - $\sigma(e^+e^- \rightarrow \text{hadrons})$  below 1 GeV dominated by  $e^+e^- \rightarrow \pi^+\pi^-$  and  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$





- New measurement of  $\sigma(e^+e^- \rightarrow \pi^+\pi^-\pi^0)$  using  $\sigma(e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma_{ISR})$  and the beam-energy constraint: see [L. Corona at Moriond 2024](#)
- Achieved accuracy of 2.2%, *moves the global fit up*
  - Dominant systematics:  $\pi^0$  eff, PROKHARA MC generator (no NNLO ISR)
  - Measurements of  $\sigma(e^+e^- \rightarrow \pi^+\pi^-)$  and others will come next
- **Tau EDM:** use spin correlation in  $e^+e^- \rightarrow \tau^+\tau^-$ , probe  $\gamma\tau\tau$  vertex vs CP reversal
  - Belle result is the world best (precision  $\sim 0.6 \times 10^{-17} e\cdot\text{cm}$ ), 20 orders above SM
  - Belle II can improve further

- What if we get a **polarised** electron beam?
  - ~70% polarisation can be a realistic target, without disruption the core physics programme (= no luminosity loss)
- **Electroweak measurements:** asymmetry in cross-sections with left- vs right-handed electrons
  - measure the neutral-current vector coupling or  $\sin \theta_W$  at 10 GeV
- Access to **g-2 (tau)** down to the SM value, and improved EDM
  - changing the beam polarisation direction is required
- Improvement in **tau Michel parameters** measurement
- Reduced backgrounds in  $\tau \rightarrow \ell \gamma$  search: SM backgrounds gets modified angular distribution
  - but what if the LFV process also gets modified? = access to helicity structure of new physics
- **Feasibility studies ongoing.**

# Direct searches

Projections

visible dark photon

scalar in  $B \rightarrow KS$ 

- A plethora of searches done, ongoing or planned:
  - Axion-like particles with  $e^+e^- \rightarrow a\gamma$  and  $a \rightarrow \gamma\gamma$ , or  $e^+e^- \rightarrow ae^+e^-$
  - Dark photons in various signatures:  $e^+e^- \rightarrow \gamma X$ ,  $e^+e^- \rightarrow \mu^+\mu^-X$ , ...  
- recent search
  - Z' in parameter space relevant for muon g-2
  - Dark matter candidates: long-lived particles, scalars in  $B \rightarrow KS$ ...
  - Dilepton resonance: recent dimuon search, probing ATOMKI anomaly in dielectron
- **Expected world-best sensitivity for many signatures below 10 GeV**
- Searches that rely on missing energy depend severely on the detector performance
  - Ensuring the **hermeticity**: a small inefficiency in one subsystem can severely impact the reach
  - Cosmic-ray veto performance



- There are many classes of "anomalies" where Belle II can contribute
  - or create new anomalies!
- Eagerly waiting for more data
- New ideas to improve the effective sensitivity even with existing data
-