
# EWP B decays with missing energy: status and future potential of Belle II analyses

Elisa Manoni - INFN Perugia

Belle II Physics week October 8<sup>th</sup>, 2025 KEK



#### Preface


- Exhaustive overview of the results in Gaetano and Lorenz's talk
- I will discuss the conventional steps of EWP B decays with missing energy (EWP+missE), with some emphasis on systematic uncertainties
- I will highlight the commonalities and distinctive features of the various works, selected based on personal basis.
- I will mainly focus on tagged analyses, inclusive method discussed in detailed by <u>Sasha</u> yesterday

# Introduction



#### Theoretical motivations

EWP B decays with missing energy (EWP+missE) as probe for new Physics



$$\int_{\Re \mathbb{R}^{n}}^{|\mathcal{A}=6|} = \sum_{i} \frac{C_{i}^{(6)}}{\Lambda^{2}} O_{i}^{(6)} [\varphi]$$
 in general violate all the accidental symmetries and properties of the SM

We can expect large effects in rare or forbidden processes!

**Precision tests** of forbidden or suppressed processes in the SM are powerful probes of physics Beyond the Standard Model.

>> Flavour Physics! <<

$$\frac{\int C}{C_{SH}} \sim \frac{C}{\lambda_{SM}} \frac{v^2}{\Lambda^2} < 10\%$$
for c=1
$$1 > \sqrt{\frac{0.8}{\lambda_{SM}}} \sqrt{10\%} \sim \frac{0.8}{\sqrt{\lambda_{SM}}} \sqrt{10\%}$$

Measuring this precisely puts strong constraints on the EFT combination c/ $\Lambda^2$ , the better the smallest  $\lambda_{SM}$  is.

Exp

TH

- Very large statistics
- Small backgrounds and systematics
- Good control over the SM prediction:
  - **SM inputs** (CKM matrix elements)
  - QCD matrix elements (form factors)
  - control over the possible long-distance contributions

In rarity lies power


D. Marzocca


# EWP+missE searches @ Belle II (I)

- Belle II is well suited to study missing energy modes:
  - constrained kinematics and detector hermeticity good missing momentum reconstruction
  - low average multiplicity neutral reconstruction

- Experimental challenges:
  - low (or zero) branching fraction with large background
  - apart from  $b \rightarrow s \tau \ell$  case, signal is non-peaking: multiple undetected neutrinos, no narrow signal structure


simulated B+→K+νν event






# EWP+missE searches @ Belle II (I)

Three main categories of EWP+missE modes studies @ Belle II:




# Tagging techniques




- So far, most of the analyses have been performed using HAD Full Event Interpretation (FEI)
- First STA results to come next year
- When applicable, ITA has proved to be more sensitive than HTA

# Typical analysis flow


#### Reconstruction and basic selection




#### Data-MC corrections




#### Background suppression



#### **Validation**



#### Signal extraction and Statistical interpretation



# Branching Fraction as main measurable

In the SM, EWP+missE modes are rare or forbidden.

The main measurable is the branching fraction, on which we usually set upper limits:

$$\mathcal{B} = \frac{N_{sig}}{N_{B\bar{B}} \cdot \varepsilon_{sig}}$$

- $\bullet$   $N_{sig}$ : from sample-composition fit or cut-and-count
- $\bullet$   $\varepsilon_{sig}$ : from signal MC
- $N_{B\bar{B}}$ : from external inputs (f<sub>+-</sub>/f<sub>00</sub> and  $\Upsilon(4S)$  counting)


Both  $N_{sig}$  and  $\varepsilon_{sig}$  's estimates require corrections to account for data/simulation discrepancies and bring the dominant systematic uncertainties

# Uncertainty balance

• The balance between statistical and systematic uncertainty depends on signal extraction strategy and working point (= background contamination), same examples:


Analysis	Signal extraction strategy	Typical dominant syst. unc.	BF uncertainty balance	Measured BF uncertainty	Sample
B+→K+νν ITA		Knowledge of background	Stat = Syst	± 0.7 x 10 <sup>-5</sup>	
B+→K+νν HTA	Binned fit	normalization, limited size of simulated sample, syst for specific backgrounds/neutral	Stat dominates but syst approaching	± 1.1 x 10 <sup>-5</sup>	Belle II
B <sup>0</sup> →K* <sup>0</sup> ττ HTA		energy corrections		± 1.0 x 10 <sup>-3</sup>	
B+→K+ττ HTA	Cut-and-count	Knowledge of background normalization	Stat dominates but syst approaching	$^{+3.7}_{-3.3} \times 10^{-4}$	BELLE
B <sup>0</sup> →K* <sup>0</sup> τℓ HTA	Unbinned fit	Knowledge of signal efficiency and signal PDF parameters	Negligible systematic uncertainty	~ ± (1.5-2.8) x 10 <sup>-5</sup>	Belle II

# Analysis dissection




# B<sub>tag</sub> reconstruction and efficiency calibration from FEI

 B<sub>tag</sub> reconstructed using "FEI" (see Karim's <u>talk</u> for more details)



- Goodness of B<sub>tag</sub> si quantified by the "FEI" signal probability
- Different  $\mathcal{P}_{FEI}$  working points recommended:
  - for HTA:  $>10^{-2}$ ,  $>10^{-3}$  (most used)
- The choice is analysis dependent: @ analysis design stage, it's worth checking different values
- Corrections are given on a Btag mode-dependent basis, with uncertainty of the order of 5% (usually, negligible impact on BF systematic error balance)

# Alternative FEI calibration: signal embedding (I)




Alternative FEI calibration technique: signal embedding (now official basf2 tool)

- Reconstruct  $B \rightarrow J/\psi(\mu\mu)K^{(*)}$  events (large branching ratio, experimentally very clean)
- Remove J/ $\psi(\mu\mu)$ K<sup>(\*)</sup> from reconstructed objects and replace with a signal B candidate from simulated signal events (both in data and MC)
- ROE (or B<sub>tag</sub>+extra particles) is untouched

# Alternative FEI calibration: signal embedding (II)

- Before using it:
  - check that the embedded sample is a good control sample for the signal (always do that when you choose a sideband)
  - check Data/MC agreement for selection variable
- Data/MC efficiency ratio used as tag efficiency correction
  - usually consistent with official FEI corrections



- CONTRA: low statistics
- PRO: analysis-specific control sample, can be use to validate signal extraction variable and full signal efficiency (and assign systematic uncertainty→ has an impact)
  - this method is indeed used also in ITA for full signal efficiency validation

# B<sub>sig</sub> reconstruction: strange mesons

- Also signal side reconstruction can be done exclusively or inclusively (or inbetween)
  - semi-exclusive reconstruction performed in  $B \rightarrow X_s \nu \nu$  search
- Strange meson reconstruction (K, K<sub>s</sub>, K\*0), starts from tracks+PID requirements as per performance group recommendation (more on <u>Bianca</u>, <u>Eldar</u> and <u>Ale</u>'s talks tomorrow)


 $B^+ \rightarrow K^+ \tau \tau$ 

 Usually, multiple PID working points are tested and the associated systematic uncertainty is sub-dominant or negligible from the ACC, CDC, and TOF for Belle. For Belle II, information from all subdetectors except the PXD and SVD is used, resulting in a kaon identification efficiency of 85% (88% for Belle) for a pion fake rate of 5% (8% for Belle) and a pion identification efficiency of 90% (91% for Belle) for a kaon fake rate of 7% (6% for Belle) at the particle identification working points  $\mathcal{P}_K, \mathcal{P}_{\pi} > 0.6$ . For Belle, electrons are identified using the information from the ECL, CDC and ACC, and information from the KLM only is used for muon identification. Using the likelihood ratio requirement  $\mathcal{P}_e(\mathcal{P}_{\mu}) > 0.9$ , the lepton identification has an efficiency of 92% (89%) and a pion fake rate of 0.3% (1.4%) for electrons (muons with momentum larger than 0.6 GeV/c). For Belle II, the electron identification uses a boosted-decision-tree (BDT) classifier trained with information from all sub-detectors except the PXD and SVD and the muon identification uses information from all sub-detectors except the PXD and SVD. At the particle identification working point  $\mathcal{P}_e(\mathcal{P}_{\mu}) > 0.9$ , the electron (muon) identification has an efficiency of 86% (89%) and a pion mis-identification rate of 0.4% (7%).

# B<sub>sig</sub> reconstruction: leptons (1)

In most of the analyses,  $\tau$  is reconstructed exclusively:

- $\tau \rightarrow \ell$ ,  $\ell = e$ ,  $\mu$  always included
- ullet in some cases,  $\pi$  and  $\rho$  modes are also considered





Even if channel with hadrons are less pure than leptonic modes, ~35% reduction on  $B^0 \rightarrow K^{*0}\tau\tau$  BF uncertainty when adding them (analysis dependent)

# B<sub>sig</sub> reconstruction: leptons (II)

In  $B \rightarrow K^{*0}\tau \ell$  search,  $\tau$  is reconstructed with an inclusive technique:

- $\tau \rightarrow$  1-prong track without particle ID, allow for up to 2 extra-tracks to includes also  $\tau \rightarrow$  3-prong decays
- comparison with exclusive reconstruction performed:
  - slightly higher efficiency in inclusive approach
  - τ invariant mass resolution essentially unchanged






# Treatment of extra-particles

- ullet In tagged analyses, events with extra-tracks or neutral extra-mesons such as  $\pi^0$  or  $K_s$  are usually rejected
- Extra-photons total energy is one of the most discriminating variables in missing energy analyses

$$E_{\text{ECL}} = \sum_{i} E_{i}$$
 $extra-\gamma$ 

Ideally, for correctly reconstructed signal events, no extra activity in the calorimeter is expected.


Tails in the signal distribution are due to mis-reconstruction effects and ECL energy resolution.



Larger ECL deposits are expected from background events. Peak at zero due to energy threshold adopted in extra photon selection

#### E<sub>ECL</sub> calibration

 Extra-photons originates from real photons, beam backgrounds, ECL clusters from tracks for which cluster-track mating failed





Data/MC agreement in E<sub>ECL</sub> depends on the extra-photon selection and on the MC

campaign. Ad hoc correction may be needed:

Event-by-event correction based on photon multiplicity distribution in some sideband and apply to EECL in the single region.


Remaining discrepancy → systematic uncertainty (has an impact)






# Strategies for background suppression


- After signal side reconstruction and requirements on ROE/extra-particles, the final background suppression stage is implemented
- Identify discriminant and well-described variables
  - E<sub>ECL</sub>, P<sub>miss</sub>, kinematic properties of signal side particles
  - Variables targeting specific backgrounds (see next slide)
  - Event shape variables to suppress continuum background
- Combined in a multivariate tool or used in a cut-andcount analysis.






# A remark on dominant common backgrounds

- Dominant backgrounds are usually from semileptonic B decays in which the signal K(\*) actually come from a D(\*)
- The invariant mass of the signal  $K^{(*)}$  and another track in the event (e.g. a signal side lepton in b $\rightarrow$ s  $\tau\ell/\tau\tau$ ) can be used to reject background







# An example of cut-and-count analysis: $B^+ \rightarrow K^+ \tau \tau$

- $\bullet$   $\tau$  reconstructed in leptonic modes
- m(Kℓ) cut to reject most of the background
- Final signal region defined by  $M^2_{miss}$ ,  $\tau$  daughter momentum and  $E_{ECL}$
- Use E<sub>ECL</sub>, B<sub>tag</sub> M<sub>BC</sub> and q<sup>2</sup> sidebands to correct background estimate

 Dominant systematic uncertainty from background and FEI corrections







An example of unbinned fit analysis:  $B^O \rightarrow K_s \tau \ell$ 

Discriminant variables combined in a BDT

- In this case, we have a variable which is peaky for the signal:  $m\tau$
- Used as signal extraction variable


 Dominant systematics from calibration of BDT cut efficiency and signal PDF parameterisation






# Examples of binned fit analyses: $B^+ \rightarrow K^+ \nu \nu$ (I)

- Final background suppression via BDT, also used as signal extraction variable
- HTA: control samples used to correct the qq and generic BB normalisation before the BDT training; data/MC validation repeated in the signal region to assign systematic uncertainty
- ITA: data/MC comparison in off-resonance data is used to background normalization systematics
- In both cases, corrections for specific backgrounds are applied before the classifier training






# Example of specific background study in $B^+ \rightarrow K^+ \nu \nu$

• Study pion-enriched control sample (B+ $\rightarrow \pi$ +X instead of B+ $\rightarrow$ K+X)

- Observed data excess in q<sup>2</sup><sub>rec</sub> distribution above D threshold
- Excess fixed by increasing  $B \rightarrow D \rightarrow K_L$  component by +30%
  - derived from 3-component fit to q<sup>2</sup><sub>rec</sub>
- Procedure successfully validated on electron- and muonenriched control samples
- 10% systematic uncertainties to cover differences in scaling factor from the different sidebands



# Systematic Uncertainties for $B^+ \rightarrow K^+ \nu \nu$

IIA	
· · · · ·	
	-

Source	Impact on $\sigma_{\mu}$
1 Normalization of $B\bar{B}$ background	0.90
Normalization of continuum background	0.10
Leading $B$ -decay branching fractions	0.22
<sup>3</sup> Branching fraction for $B^+ \to K^+ K_L^0 K_L^0$	0.49
p-wave component for $B^+ \to K^+ K_{\rm S}^0 K_{\rm L}^0$	0.02
<b>4</b> Branching fraction for $B  o D^{**}$	0.42
Branching fraction for $B^+ \to K^+ n\bar{n}$	0.20
Branching fraction for $D \to K_L^0 X$	0.14
Continuum-background modeling, BDT <sub>c</sub>	0.01
Integrated luminosity	< 0.01
Number of $B\bar{B}$	0.02
Off-resonance sample normalization	0.05
Track-finding efficiency	0.20
Signal-kaon PID	0.07
Photon energy	0.08
Hadronic energy	0.37
$K_{\rm L}^0$ efficiency in ECL	0.22
Signal SM form-factors	0.02
Global signal efficiency	0.03
2 Simulated-sample size	0.52

HTA	
Source	Impact on $\sigma_{\mu}$
1 Normalization of $B\bar{B}$ background	0.91
4 Normalization of continuum background	0.58
Leading <i>B</i> -decay branching fractions	0.10
Branching fraction for $B^+ \to K^+ K_L^0 K_L^0$	0.20
Branching fraction for $B \to D^{**}$	< 0.01
Branching fraction for $B^+ \to K^+ n\bar{n}$	0.05
Branching fraction for $D \to K_L^0 X$	0.03
Continuum-background modeling, BDT <sub>c</sub>	0.29
Number of $B\bar{B}$	0.07
Track finding efficiency	0.01
Signal-kaon PID	< 0.01
<sup>2</sup> Extra-photon multiplicity	0.61
$K_{\rm L}^0$ efficiency	0.31
Signal SM form-factors	0.06
Signal efficiency	0.42
3 Simulated-sample size	0.60

# Pre box-opening consistency checks

Once the analysis is frozen, before looking the full signal region, perform test on portions of the dataset or on control samples to see that you get what you expect

1) Fit to the **same-flavor** data  $\rightarrow$  BR, p-value, pulls


 $B^0 \rightarrow K^{*0} \tau \tau$ 

- 2) Partial fit
  - a) signal region restricted to the lowest sensitive BDT bins with all signal categories  $\rightarrow$  BR, p-value, pulls
- 3) Blind fit on **half-split proc vs prompt**, where only the uncertainty and the μ differences between the two samples are compared

 $B^+ \rightarrow K^+ \nu \nu$ 


- 4) Full fit to the signal region keeping all results blinded expulls of nuisances
- 5) Partial fit
  - b) signal region restricted to  $\pi\pi$  category  $\rightarrow$  BR, p-value, pulls


Divide data sample into pairs of statistically independent datasets, according to various features





- Good stability for all splittings for both analyses
- Excellent agreement when splitting ITA sample according to lepton multiplicity (probing "semileptonic tag" vs "hadronic tag")
- Tension in "Sum(charges)" for ITA consistent with statistical fluctuation

### Gran-summary









# What's next

# Improve sensitivity with Bellell data

- $b \rightarrow s \nu \nu$ : unique to Belle II, corroborate evidence improving precision with
  - ITA with additional K<sup>(\*)</sup> channels
  - additional tagging techniques (STA to improve precision, HTA to study q<sup>2</sup> spectrum with better resolution)
  - use of Run2 dataset (and what will come next)
- b→sττ:
  - most stringent limit from  $B^0 \rightarrow K^{*0}\tau\tau$  LHCb measurement (Abhijit's talk)
  - many handles to improve Belle II  $B^+ \to K^+ \tau \tau$  measurement: additional  $\tau$  channels, semileptonic and inclusive tag, further optimisation of analysis strategy when adding less pure modes
- b→d effort also started

# Future perspective in systematics

We foreseen improvements on systematics related to:

- background normalisations →generally, statistical in origin
- specific background modelling
- detectors performances (improvements in particle reconstruction and identification algorithms, usage of MCrd)
- simulated sample size → larger MC samples can be produced in a smart way (save to disk only skimmed events)

Source	Impact on $\sigma_{\mu}$
Normalization of $B\bar{B}$ background	0.90
Normalization of continuum background	0.10
Leading B-decay branching fractions	0.22
Branching fraction for $B^+ \to K^+ K_{\rm L}^0 K_{\rm L}^0$	0.49
p-wave component for $B^+ \to K^+ K_{\rm S}^0 K_{\rm L}^0$	0.02
Branching fraction for $B \to D^{**}$	0.42
Branching fraction for $B^+ \to K^+ n\bar{n}$	0.20
Branching fraction for $D \to K_L^0 X$	0.14
Continuum-background modeling, BDT <sub>c</sub>	0.01
Integrated luminosity	< 0.01
Number of $B\bar{B}$	0.02
Off-resonance sample normalization	0.05
Track-finding efficiency	0.20
Signal-kaon PID	0.07
Photon energy	0.08
Hadronic energy	0.37
$K_{\rm L}^0$ efficiency in ECL	0.22
Signal SM form-factors	0.02
Global signal efficiency	0.03
Simulated-sample size	0.52

# Summary

- EWP+missE are excellent probe for New Physics
- Analysis are challenging, background validation is a key point
- Systematics starts to be important for all analyses, handles to improve them
- A lot of work ahead for Belle II to continue producing world-leading results
  - combination of tagging technique
  - improvements in reconstruction, MC modelling, and analysis technique are within reach



# Extra slides

### References

	Analysis	Journal reference
b→s νν	B+→K+νν [doc] B→Xsνν [doc]	PRD 109 (2024) 11, 112006, to be submitted to PRL
b→s ττ	$B^{+} \rightarrow K^{+} \tau \tau \ [\underline{doc}]$ $B^{0} \rightarrow K^{*0} \tau \tau [\underline{doc}]$	to be submitted to PRL, PRL 135 (2025), 151801
b→s τℓ	$B^{0} \rightarrow K_{S} \tau \ell \text{ [doc]}$ $B^{0} \rightarrow K^{*0} \tau \ell \text{ [doc]}$	PRL 135 (2025) 4, 041801 JHEP 08 (2025) 184

#### $B^+ \rightarrow K^+ \nu \nu ITA$

		<u> </u>		
Source	Correction	Uncertainty type, parameters	Uncertainty size	Impact on $\sigma_{\mu}$
Normalization of $B\bar{B}$ background		Global, 2	50%	0.90
Normalization of continuum background		Global, 5	50%	0.10
Leading <i>B</i> -decay branching fractions		Shape, 6	O(1%)	0.22
Branching fraction for $B^+ \to K^+ K_{\rm L}^0 K_{\rm L}^0$	$q^2$ dependent $O(100\%)$	Shape, 1	20%	0.49
p-wave component for $B^+ \to K^+ K_S^0 K_L^0$	$q^2$ dependent $O(100\%)$	Shape, 1	30%	0.02
Branching fraction for $B \to D^{**}$	7	Shape, 1	50%	0.42
Branching fraction for $B^+ \to K^+ n\bar{n}$	$q^2$ dependent $O(100\%)$	Shape, 1	100%	0.20
Branching fraction for $D \to K_L^0 X$	+30%	Shape, 1	10%	0.14
Continuum-background modeling, BDT <sub>c</sub>	Multivariate $O(10\%)$	Shape, 1	100% of correction	0.01
Integrated luminosity		Global, 1	1%	< 0.01
Number of $B\bar{B}$		Global, 1	1.5%	0.02
Off-resonance sample normalization		Global, 1	5%	0.05
Track-finding efficiency		Shape, 1	0.3%	0.20
Signal-kaon PID	$p, \theta$ dependent $O(10-100\%)$	Shape, 7	O(1%)	0.07
Photon energy		Shape, 1	0.5%	0.08
Hadronic energy	-10%	Shape, 1	10%	0.37
$K_{\rm L}^0$ efficiency in ECL	-17%	Shape, 1	8.5%	0.22
Signal SM form-factors	$q^2$ dependent $O(1\%)$	Shape, 3	O(1%)	0.02
Global signal efficiency		Global, 1	3%	0.03
Simulated-sample size		Shape, 156	O(1%)	0.52

### $B^+ \rightarrow K^+ \nu \nu HTA$

Source	Correction	Uncertainty type, parameters	Uncertainty size	Impact on $\sigma_{\mu}$
Normalization of $B\bar{B}$ background		Global, 1	30%	0.91
Normalization of continuum background		Global, 2	50%	0.58
Leading B-decay branching fractions		Shape, 3	O(1%)	0.10
Branching fraction for $B^+ \to K^+ K_{\rm L}^0 K_{\rm L}^0$	$q^2$ dependent $O(100\%)$	Shape, 1	20%	0.20
Branching fraction for $B \to D^{**}$		Shape, 1	50%	< 0.01
Branching fraction for $B^+ \to K^+ n\bar{n}$	$q^2$ dependent $O(100\%)$	Shape, 1	100%	0.05
Branching fraction for $D \to K_L^0 X$	+30%	Shape, 1	10%	0.03
Continuum-background modeling, BDT <sub>c</sub>	Multivariate $O(10\%)$	Shape, 1	100% of correction	0.29
Number of $B\bar{B}$		Global, 1	1.5%	0.07
Track finding efficiency		Global, 1	0.3%	0.01
Signal-kaon PID	p, $\theta$ dependent $O(10-100\%)$	Shape, 3	O(1%)	< 0.01
Extra-photon multiplicity	$n_{\gamma \text{extra}}$ dependent $O(20\%)$	Shape, 1	O(20%)	0.61
$K_{\rm L}^0$ efficiency		Shape, 1	17%	0.31
Signal SM form-factors	$q^2$ dependent $O(1\%)$	Shape, 3	O(1%)	0.06
Signal efficiency		Shape, 6	16%	0.42
Simulated-sample size		Shape, 18	O(1%)	0.60

# $B \rightarrow X_s \nu \nu$

Source	Impact on $\sigma_{\mathcal{B}}$ [10 <sup>-1</sup>
Simulated-sample size	6.0
Background normalization	5.7
Branching ratio of major $B$ meson decay	2.3
Non-resonant $X_s \nu \bar{\nu}$ generation point	2.1
$\mathcal{O}_{\mathrm{BDT}}$ selection efficiency	2.0
Photon multiplicity correction	1.8
$q\bar{q}$ background efficiency	1.8
Other subdominant contributions	3.0
Total systematic sources	11.7

Source	Impact on $\mathcal{B} \times 10^{-3}$
$B \to D^{**} \ell / \tau \nu$ branching fractions	0.29
Simulated sample size	0.27
$q\bar{q}$ normalization	0.18
ROE cluster multiplicity	0.17
$\pi$ and $K$ ID	0.14
B decay branching fraction	0.11
Combinatorial $B\bar{B}$ normalization	0.09
Signal and peaking $B^0\bar{B}^0$ normalization	0.07
Lepton ID	0.04
$\pi^0$ efficiency	0.03
$f_{00}$	0.01
$N_{\Upsilon(4S)}$	0.01
$D \to K_L^0$ decays	0.01
Signal form factors	0.01
Luminosity	< 0.01
Total systematics	0.52
Statistics	0.86

Signal category	$\varepsilon \times 10^5$	$Bar{B}$	$qar{q}$
$\ell\ell$	$4.0 \pm 0.6$	275	39
$\pi\ell$	$7.6 \pm 1.1$	1058	230
ho	$15.5 \pm 2.2$	3279	845
$\pi\pi$	$4.0 \pm 0.6$	1077	424

Sources	Impact on BF (Belle)	Impact on BF (Belle II)
Expected bkg yield	±2.5 events	±1.2 events
FEI scale factor	10.1%	12.6%
Simulated sample size	3.3%	3.5%
PID correction	1.0%	1.6%
pi0 veto	1.9%	2.9%
Tracking efficiency	1.1%	0.8%
Signal decay model	3.5%	4.3%
f+-	+1.4 % -2.1 %	+1.4 % -2.1 %
Number of BBbar pairs	1.4%	1.6%

Source (%)	$K_S^0 au^+\mu^-$	$K_S^0  au^- \mu^+$	$K_S^0  au^+ e^-$	$K_S^0  au^- e^+$
BDT selection	17.1	17.5	16.6	19.2
Signal PDF	15.7	15.7	15.7	15.7
$B_{ m tag}$ calibration	3.7	3.7	3.7	3.7
Lepton identification	0.3	0.3	0.5	0.5
$\tau$ daughter reconstruction	5.3	6.5	6.1	6.4
Linearity	1.6	1.4	0.8	1.4
Number of $B\bar{B}$ pairs	1.1	1.1	1.1	1.1
$f_{+-}/f_{00}$	2.3	2.3	2.3	2.3
$K_S^0,  \tau,  \rho,  \pi^0$ branching fractions	0.7	0.7	0.7	0.7
Total	24	24	23	25

			$\mathcal{B}(10^{-5})$	
Channels	$\epsilon(10^{-4})$	$N_{ m sig}$	Central value	UL
$B^0 \to K_S^0 \tau^+ \mu^-$	1.7	$-1.8 \pm 3.0$	$-1.0 \pm 1.6 \pm 0.2$	1.1
$B^0 \to K_S^0 \tau^- \mu^+$	2.1	$2.6 \pm 3.5$	$1.1 \pm 1.6 \pm 0.3$	3.6
$B^0 \to K_S^0 \tau^+ e^-$	2.0	$-1.2\pm2.4$	$-0.5 \pm 1.1 \pm 0.1$	1.5
$B^0 \to K_S^0 \tau^- e^+$	2.1	$-2.9 \pm 2.0$	$-1.2 \pm 0.9 \pm 0.3$	0.8

# $B^0 \rightarrow K^{*0} \tau \ell$

Source	Belle	Belle II	
FEI efficiency	4.9%(OSe), 4.9%(SSe),	6.2%(OSe), 6.1%(SSe),	
1 Li efficiency	$4.9\%(OS\mu), 4.9\%(SS\mu)$	$6.1\%(OS\mu), 6.2\%(SS\mu)$	
Lepton ID efficiency	2.0%(OSe), 2.4%(SSe),	0.7%(OSe), 1.1%(SSe),	
	$2.2\%(OS\mu), 2.2\%(SS\mu)$	$0.7\%(OS\mu), 0.6\%(SS\mu)$	
Hadron ID efficiency	1.9%(OSe), 2.0%(SSe),	3.7%(OSe), 3.7%(SSe),	
	$1.9\%(OS\mu), 2.0\%(SS\mu)$	$3.6\%(OS\mu), 3.7\%(SS\mu)$	
BDT efficiency	27%(OSe), 21%(SSe),	29%(OSe), 31%(SSe),	
Tracking officioner	$18\%(OS\mu), 23\%(SS\mu)$ 1.4%	$34\%(OS\mu), 31\%(SS\mu)$ 1.1%	
Tracking efficiency	27.6%(OSe),	29.8% (OSe),	
	21.8%(SSe),	31.8%(SSe),	
Total efficiency	$18.9\%(OS\mu),$	$34.7\%(OS\mu),$	
	$23.7\%(SS\mu)$	$31.7\%(SS\mu)$	
Signal mean $\mu$	0.1%	0.2%	
Signal width $\lambda$	21%	59%	
Background shape exp $(\times 10^{-5})$	$0.1(OSe),0.1(SSe),0.1(OS\mu),0.6(SS\mu)$		
Background shape obs $(\times 10^{-5})$	$0.1(OSe),0.3(SSe),0.1(OS\mu),0.1(SS\mu)$		
$B\overline{B}$ pairs	1.4%	1.6%	
$f^{00}$	0.8%		
$\mathcal{B}r(K^{*0} \to K^+\pi^-)$	< 0.05%		
MC statistics	< 0.0	0005%	

Decay	$\mathcal{B}^{\text{fit}} (\times 10^{-5})$	$\mathcal{B}_{\text{obs(exp)}}^{\text{UL}} \ (\times 10^{-5})$
$OSe: B^0 \to K^{*0}\tau^+e^-$	$-0.24 \pm 1.46$	2.9 (2.8)
$SSe: B^0 \to K^{*0}\tau^-e^+$	$1.17 \pm 2.77$	6.4 (4.4)
$OS\mu: B^0 \to K^{*0}\tau^+\mu^-$	$1.07 \pm 1.80$	4.2(3.0)
$SS\mu: B^0 \to K^{*0}\tau^-\mu^+$	$0.48 \pm 2.61$	5.6(5.5)

		OSe	SSe	$OS\mu$	$SS\mu$
Belle	$N_{ m sig}$	$-0.6 \pm 3.3$	$2.2 \pm 5.2$	$2.8 \pm 4.7$	$0.6 \pm 3.2$
			$140.0 \pm 12.7$		
Belle II	$N_{ m sig}$	$-0.5 \pm 2.8$	$1.7 \pm 3.9$	$1.6 \pm 2.7$	$0.6 \pm 3.3$
	$N_{ m bkg}$	$86.5 \pm 9.8$	$120.1\pm11.8$	$109.1\pm10.9$	$198.4 \pm 14.5$

#### $B \rightarrow X_s \nu \nu$

- A special case:  $B \rightarrow X_s \nu \nu$  with  $X_s = sum$ of-exclusive modes
- 30 modes, ~93% of entire X₅ sample based on Belle II MC (PYTHIA8)

		$B^{0}\bar{B}^{0}$			$B^{\pm}$	
		<i>D D</i>			<i>D</i>	
K	$K_S^0$			$K^{\pm}$		
$K\pi$	$K^{\pm}\pi^{\mp}$	$K_S^0\pi^0$		$K^{\pm}\pi^0$	$K_S^0 \pi^{\pm}$	
$K2\pi$	$K^{\pm}\pi^{\mp}\pi^{0}$	$K_S^0\pi^\pm\pi^\mp$	$K^0_S\pi^0\pi^0$	$K^{\pm}\pi^{\mp}\pi^{\pm}$	$K_S^0 \pi^{\pm} \pi^0$	$K^{\pm}\pi^0\pi^0$
$K3\pi$	$K^{\pm}\pi^{\mp}\pi^{\pm}\pi^{\mp}$	$K_S^0 \pi^{\pm} \pi^{\mp} \pi^0$	$K^{\pm}\pi^{\mp}\pi^{0}\pi^{0}$	$K^{\pm}\pi^{\mp}\pi^{\pm}\pi^{0}$	$K_S^0 \pi^{\pm} \pi^{\mp} \pi^{\pm}$	$K_S^0\pi^\pm\pi^0\pi^0$
$K4\pi$	$K^{\pm}\pi^{\mp}\pi^{\pm}\pi^{\mp}\pi$	${}^{0}K_{S}^{0}\pi^{\pm}\pi^{\mp}\pi^{\pm}\pi^{\pm}$	$\mp K_S^0 \pi^{\pm} \pi^{\mp} \pi^0 \pi^0$	$K^{\pm}\pi^{\mp}\pi^{\pm}\pi^{\mp}\pi$	$\pm K_S^0 \pi^{\pm} \pi^{\mp} \pi^{\pm} \pi^0$	$0 K^{\pm}\pi^{\mp}\pi^{\pm}\pi^{0}\pi^{0}$
3K	$K^{\pm}K^{\mp}K^0_S$			$K^{\pm}K^{\mp}K^{\pm}$		
$3K\pi$	$K^{\pm}K^{\mp}K^{\pm}\pi^{\mp}$	$K^{\pm}K^{\mp}K^0_S\pi^0$		$K^{\pm}K^{\mp}K^{\pm}\pi^{0}$	$K_S^0 K^{\pm} K^{\mp} \pi^{\pm}$	

- Signal model: exclusive  $B \rightarrow K \nu \nu$  and  $B \rightarrow K^* \nu \nu + \text{and } B \rightarrow X_s \nu \nu$  for  $M_{Xs} > 1.1$  GeV/c<sup>2</sup> (systematic on endpoint variation, mid-importance)
  - Fermi motion model, assuming Gaussian b-quark momentum distribution.
  - Final-state hadrons fragmentation simulated via Pythia8 and corrected using  $B \rightarrow X_s \gamma$  study (negligible associated systematics)