

Recent Dark Sector results at Belle II



Giacomo De Pietro



for the Belle II collaboration




Identification of Dark Matter @ Vienna
18-22 September 2022

SuperKEKB: a new Intensity Frontier machine

SuperKEKB is a **super** B-factory located at KEK (Tsukuba, Japan)


It's an asymmetric e^+e^- collider operating mainly at **10.58 GeV** ($\Upsilon(4S)$, but possible runs from $\Upsilon(2S)$ to $\Upsilon(6S)$)



I (A): $\sim 1.6/1.2$


β_y^* (mm): $\sim 5.9/5.9$

nano-beam
scheme



I (A): $\sim 2.8/2.0$

β_y^* (mm): $\sim 0.27/0.3$




30x peak luminosity:
 $6 \cdot 10^{35} \text{ cm}^{-2} \text{ s}^{-1}$

SuperKEKB: a new Intensity Frontier machine

SuperKEKB is
located at KEK



It's an asymmetric
operating mainly
($\Upsilon(4S)$), but possible run

KEKB



$I(A): \sim 1.6/1.2$

$\beta_y^* (mm): \sim 5.9/5.9$



Peak luminosity:
 $10^{35} \text{ cm}^{-2} \text{ s}^{-1}$

Belle II: a new Intensity Frontier detector

Electromagnetic Calorimeter (ECL):

CsI(Tl) crystals, waveform sampling to measure time, energy, and pulse-shape.

K_L and muon detector (KLM):

Resistive Plate Counters (RPC) (outer barrel)
Scintillator + WLSF + MPPC (endcaps, inner barrel)

Magnet:

1.5 T superconducting

Trigger:

Hardware: < 30 kHz
Software: < 10 kHz

Vertex detectors (VXD):


2 layer DEPFET pixel detectors (PXD)
4 layer double-sided silicon strip detectors (SVD)

Central drift chamber (CDC):

He(50%):C₂H₆ (50%), small cells,
fast electronics


Particle Identification (PID):

Time-Of-Propagation counter (TOP) (barrel)
Aerogel Ring-Imaging Cherenkov Counter (ARICH) (FWD)



SuperKEKB and Belle II operations


First collisions: 26th April 2018



Collected 0.5 fb^{-1} in 2018

Collected about 424 fb^{-1} since 2019:

- 363 fb^{-1} at the $\Upsilon(4S)$ resonance
- 61 fb^{-1} off-resonance



Goal: integrate up to 50 ab^{-1} in a decade!


Dark Matter

It is “dark”.


It exists...

A lot of experimental techniques
to probe the Dark Matter existence:

- **production at colliders.**



Dark Matter coupling to Standard Model



Different possible portals between **Dark Matter** and **Standard Model** depending on the **dark mediator X**:

Vector portal $\rightarrow$ Dark Photon / Z'

Scalar portal $\rightarrow$ Dark Higgs / Dark Scalar

Pseudoscalar portal $\rightarrow$ Axion-Like Particles

Neutrino portal $\rightarrow$ Sterile Neutrinos



$\tau^\pm \rightarrow (e^\pm / \mu^\pm) \alpha ; \alpha \rightarrow \text{invisible}$

Can enter from NP models such as light ALP

Best upper limits on $B(\tau \rightarrow l \alpha) / B(\tau \rightarrow l \nu \bar{\nu})$ from ARGUS (476 pb⁻¹, *Z. Phys. C* 68 (1995) 25)

From phenomenology: consistency of $B(\tau \rightarrow l \nu \bar{\nu})$ with SM predictions

With current data, Belle II can already set more stringent limits



$\tau^\pm \rightarrow (e^\pm / \mu^\pm) \alpha$ - Reconstruction

Dataset: 62.8 fb⁻¹


Split event in two emispheres across thrust axis

Require exactly 4 tracks:



- 1 lepton on signal side
- 3 pions on tag side
- veto neutrals (γ , π^0) for reducing hadronic background

$\tau \rightarrow l \alpha$ events are indistinguishable from $\tau \rightarrow l \nu \bar{\nu}$ from (irreducible background)

Reducible backgrounds ($q\bar{q}$, l^+l^- , $l^+l^-l^+l^-$, $\tau^+\tau^-$ with $\tau \rightarrow \pi\nu$) suppressed with “safe” selection cuts



$\tau^\pm \rightarrow (e^\pm / \mu^\pm) \alpha$ – Data and MC spectra





$\tau \rightarrow l \alpha$ events shown normalised to $B(\tau \rightarrow l \alpha) = 5\%$

$$x_\ell \equiv \frac{E_\ell}{m_\tau/2}$$

High purity (96% for electron channel, 92% for muon channel) with the efficiency between 9% and 17% depending on M_α

No signal observed $\rightarrow$ set 95% CL upper limits



Largest systematics from particle identification

Most stringent measurements in these channels to date

We searched for a $\tau^+ \tau^-$ resonance in $\mu^+ \mu^- \tau^+ \tau^-$ final states

$$\rightarrow M(\tau\tau) = M_{\text{recoil}}(\mu\mu)$$

We probed three different models:

Z' $L_\mu - L_\tau$ model


- JHEP 12 (2016) 106
- vector portal
- first time search in $\tau\tau$

Leptophilic dark scalar S model

- PRD 95 (2017) 075003
- Yukawa couplings
- constraints by BaBar in $S \rightarrow \mu\mu$
- first time search in $\tau\tau$

$\text{ALP} \rightarrow \tau\tau$

- JHEP 12 (2017) 044
- $C_{ee} = C_{\mu\mu} = C_{\tau\tau}$; $C_{\gamma\gamma} = C_{Z\gamma} = 0$
- Yukawa-like effective couplings
- ALP- τ coupling unconstrained



$Z' / S / ALP \rightarrow \tau^+\tau^-$ - Reconstruction

Dataset: 63.3 fb^{-1}

Require exactly 4 tracks:

- $2\mu + 2e/\mu/\pi$
- $M(4 \text{ tracks}) < 9.5 \text{ GeV}$
- allowed neutrals
- scan $M_{\text{recoil}}(\mu\mu)$

Main backgrounds:

- $\tau^+\tau^-(\gamma)$
- $q\bar{q}$
- $l^+l^-l^+l^-$ (no ISR in our simulation)
- $\mu^+\mu^-\tau^+\tau^- + e^+e^-X_{\text{had.}}$ (not simulated)

$Z' / S / \text{ALP} \rightarrow \tau^+\tau^-$ - Reconstruction

Dataset: 63.3 fb^{-1}

Require exactly 4 tracks:




- $2\mu + 2e/\mu/\pi$
- $M(4 \text{ tracks}) < 9.5 \text{ GeV}$
- allowed neutrals
- scan $M_{\text{recoil}}(\mu\mu)$

Main backgrounds:

- $\tau^+\tau^-(\gamma)$
- $q\bar{q}$
- $l^+l^-l^+l^-$ (no ISR in our simulation)
- $\mu^+\mu^-\tau^+\tau^- + e^+e^-X_{\text{had.}}$ (not simulated)

Background suppression via dedicated MLPs



→ 8 MLP ranges in $M_{\text{recoil}}(\mu\mu)$



Selection optimized for $Z' \rightarrow \tau^+\tau^-$ signal

→ achieved 99% background reduction

$Z' / S / \text{ALP} \rightarrow \tau^+\tau^-$ - Data and MC spectra




Discrepancies expected and understood due to missing features in simulation

No peaking structures in $M_{\text{recoil}}(\mu\mu)$


Signal mass resolution from 1.5 MeV to 30 MeV

No signal observed $\rightarrow$ set 90% CL upper limits




Results on $Z' \rightarrow \tau\tau$ not competitive w.r.t. other Z' searches

First constraints on S for $M_S > 6.5 \text{ GeV}$



First direct constraints on $\text{ALP} \rightarrow \tau\tau$



Summary

- ✓ Belle II will lead the field in the Dark Sector searches in the MeV-GeV mass range in the coming years
- ✓ Belle II collected about 424 fb^{-1} of collisions data
- ✓ Presented here world-leading results for searches of:
 - $\tau^\pm \rightarrow (e^\pm / \mu^\pm) \alpha$, with $\alpha \rightarrow \text{invisible}$
 - leptophilic dark scalar $S \rightarrow \tau^+\tau^-$ / ALP $\rightarrow \tau^+\tau^-$
- ✓ More results will be presented by M. Campajola in his plenary talk
 - Dark Higgsstrahlung $e^+e^- \rightarrow A'h'$, with $A' \rightarrow \mu^+\mu^-$ and h' invisible
 - Invisible Z' within the $L_\mu - L_\tau$ model



Thank you for
your attention





Backup
slides

B-factories as Intensity Frontier experiments

B-factories are dedicated experiments at e^+e^- asymmetric-energy colliders for the production of quantum coherent $B\bar{B}$ pairs.

First generation of B-factories

(collected about 1.5 ab^{-1} of integrated luminosity)



The strenghts of a B-factory are:




- constrained kinematics;
- clean environment and lower background;
- hermetic detector;
- excellent PID capabilities;
- efficienct reconstruction of neutral particles.

B-factories as Intensity Frontier experiments

B-factories are dedicated experiments at e^+e^- asymmetric-energy colliders for the production of quantum coherent $B\bar{B}$ pairs.

First generation of B-factories

(collected about 1.5 ab^{-1} of integrated luminosity)



SuperKEKB machine parameters

Parameter	KEKB Design	KEKB Achieved	SuperKEKB Design
Energy (GeV) (LER/HER)	3.5/8.0	3.5/8.0	4.0/7.0
β_y^* (mm)	10/10	5.9/5.9	0.27/0.30
β_x^* (mm)	330/330	1200/1200	32/25
ϵ_x (nm)	18/18	18/24	3.2/5.3
$\frac{\epsilon_y}{\epsilon_x}$ (%)	1	0.85/0.64	0.27/0.24
σ_y (μm)	1.9	0.94 $\xrightarrow{1/20}$	0.048/0.062
ξ_y	0.052	0.129/0.090	0.09/0.081
σ_z (mm)	4	6/7	6/5
I_{beam} (A)	2.6/1.1	1.64/1.19 $\xrightarrow{\times 2}$	3.6/2.6
$N_{bunches}$	5000	1584	2500
Luminosity ($10^{34} \text{cm}^{-2} \text{s}^{-1}$)	1.0	2.11 $\xrightarrow{\times 40}$	80


$$L = \frac{\gamma_{\pm}}{2er_e} \left(1 + \frac{\sigma_y^*}{\sigma_x^*} \right) \left(\frac{I_{\pm} \xi_{y\pm}}{\beta_y^*} \right) \left(\frac{R_L}{R_{\xi_{y\pm}}} \right)$$

Cross sections at a B-factory

Physics process	Cross section [nb]	Selection Criteria	Reference
$\Upsilon(4S)$	1.110 ± 0.008	-	[2]
$u\bar{u}(\gamma)$	1.61	-	KKMC
$d\bar{d}(\gamma)$	0.40	-	KKMC
$s\bar{s}(\gamma)$	0.38	-	KKMC
$c\bar{c}(\gamma)$	1.30	-	KKMC
$e^+e^-(\gamma)$	300 ± 3 (MC stat.)	$10^\circ < \theta_e^* < 170^\circ$, $E_e^* > 0.15$ GeV	BABAYAGA.NLO
$e^+e^-(\gamma)$	74.4	$p_e > 0.5$ GeV/c and e in ECL	-
$\gamma\gamma(\gamma)$	4.99 ± 0.05 (MC stat.)	$10^\circ < \theta_\gamma^* < 170^\circ$, $E_\gamma^* > 0.15$ GeV	BABAYAGA.NLO
$\gamma\gamma(\gamma)$	3.30	$E_\gamma > 0.5$ GeV in ECL	-
$\mu^+\mu^-(\gamma)$	1.148	-	KKMC
$\mu^+\mu^-(\gamma)$	0.831	$p_\mu > 0.5$ GeV/c in CDC	-
$\mu^+\mu^-\gamma(\gamma)$	0.242	$p_\mu > 0.5$ GeV in CDC, $\geq 1 \gamma$ ($E_\gamma > 0.5$ GeV) in ECL	-
$\tau^+\tau^-(\gamma)$	0.919	-	KKMC
$\nu\bar{\nu}(\gamma)$	0.25×10^{-3}	-	KKMC
$e^+e^-e^+e^-$	39.7 ± 0.1 (MC stat.)	$W_{\ell\ell} > 0.5$ GeV/c ²	AAFH
$e^+e^-\mu^+\mu^-$	18.9 ± 0.1 (MC stat.)	$W_{\ell\ell} > 0.5$ GeV/c ²	AAFH

E. Kou, P. Urquijo et al.,
arXiv:1808.10567


Electromagnetic Calorimeter (ECL)



In barrel ECL, Belle II has **no projective cracks in ϕ** w.r.t. BaBar:


→ more hermetic

→ more efficient




Electromagnetic Calorimeter (ECL)


Energy resolution in Belle II barrel:



Material budget in front of ECL:



A rule of thumb...



The masses of the mediator and of the DM candidates lead to **different type of searches**.



ALP $\rightarrow \gamma\gamma$ - Model

Axion-Like Particles (ALPs) are pseudoscalar particles (a) that couple to bosons.

Unlike QCD Axions, ALPs have no relation between mass and coupling.

Belle II focused on the **coupling to photons**:

$$\mathcal{L} \supset -\frac{g_{a\gamma\gamma}}{4} a F_{\mu\nu} \tilde{F}^{\mu\nu} \quad \tau_a \sim 1/g_{a\gamma\gamma}^2 m_a^3$$




Investigating the photon
coupling $g_{a\gamma\gamma}$ in ALP-strahlung


First search at B-factories

ALP $\rightarrow \gamma\gamma$ – Reconstruction & Data/MC spectra



Select events with no charged tracks consisting of 3 isolated photons with a total invariant mass consistent with $s^{1/2}$.

Search strategy optimized to maximize ALP sensitivity:

- low ALP mass $\rightarrow$ **diphoton mass spectrum**; 
- high ALP mass $\rightarrow$ **recoil mass spectrum**. 



PRL 125 (2020) 161806



ALP $\rightarrow \gamma\gamma$ - Results

Search conducted with 445 pb⁻¹ of **2018 pilot run data**:

- 500 fits in sliding ranges with steps of half mass resolution;
- no excess observed (largest local significance: 2.8 σ).

$$\sigma_a = \frac{g_{a\gamma\gamma}^2 \alpha_{\text{QED}}}{24} \left(1 - \frac{m_a^2}{s}\right)^3$$

