

# Z' analyses at Belle II

**Martina Laurenza (she/her)**  
for the Belle II experiment

Uppsala Universitet

✉ [martina.laurenza@physics.uu.se](mailto:martina.laurenza@physics.uu.se)




DMNet International Symposium

Padova, 26-28 September 2023

# Introduction




# Light Dark Matter at B-factories



- **Dark Matter** is one of the most compelling reasons for **New Physics**
- B-factories at  $e^+e^-$  collider can access the mass range favored by **light dark sector**  
 → **Possible sub-GeV scenario:**  
 DM weakly coupled to SM through a **light mediator X**:

1. **Vector portal**  
Dark Photons,  $Z'$  bosons
2. **Pseudo-scalar portal**  
Axion Like Particles (ALPs)
3. **Scalar portal**  
Dark higgs/Scalars
4. **Neutrino portal**  
Sterile Neutrinos


# Light Dark Matter at B-factories



- **Dark Matter** is one of the most compelling reasons for **New Physics**
- B-factories at  $e^+e^-$  collider can access the mass range favored by **light dark sector**  
 → **Possible sub-GeV scenario:**  
 DM weakly coupled to SM through a **light mediator X**:

1. Vector portal  
Dark Photons,  $Z'$  bosons
  2. Pseudo-scalar portal  
**Axion Like Particles (ALPs)**
  3. Scalar portal  
**Dark higgs/Scalars**
  4. Neutrino portal  
Sterile Neutrinos
- See L. Zani presentation


# Light Dark Matter at B-factories



- Dark Matter is one of the most compelling reasons for **New Physics**
- B-factories at  $e^+e^-$  collider can access the mass range favored by **light dark sector**
  - **Possible sub-GeV scenario:**  
DM weakly coupled to SM through a **light mediator X**:


- Vector portal**  
Dark Photons,  $Z'$  bosons
  - Pseudo-scalar portal  
Axion Like Particles (ALPs)
  - Scalar portal  
Dark higgs/Scalars
  - Neutrino portal  
Sterile Neutrinos
- This presentation

# Light Dark Matter possible signatures




- Once produced, the mediator can have three different types of decays:
  1. Invisible decays
  2. Leptonic decays
  3. Hadronic decays

# Light Dark Matter possible signatures



- Once produced, the mediator can have three different types of decays:
  - Invisible decays**
  - Leptonic decays**
  - Hadronic decays**

# Light Dark Matter possible signatures



→ In this presentation:

○ Once produced, the mediator can have three different types of decays:

1. Invisible decays:  $Z' \rightarrow \text{inv.}$
2. Leptonic decays:  $Z' \rightarrow \mu\mu$
3. Hadronic decays  $Z' \rightarrow \tau\tau$


**+ some extras**

# Dark Sector @ Belle II

\* Dedicated talk from E. Graziani

- Signature-based
- Advantages from the **low particle multiplicity** at lepton colliders + **hermetic detector**:
  - Belle II at SuperKEKB asymmetric  $e^+e^-$  collider
- running at 10.58 GeV, well-known **initial condition**
- efficient reconstruction of **neutrals**
- specific low-multiplicity **triggers (not present at Belle)**
- excellent particle identification system

**Unprecedented luminosity**  
 $4.7 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$



- Shutdown since 2022 to install two-layer pixel detector
- 424 fb<sup>-1</sup> collected to date
- Data taking resume by end of 2023



# **Z' analyses at Belle II**




# The $L_\mu - L_\tau$ model

[1] B.Shuve and I.Yavin (2014) Phys. Rev. D 89, 113004;  
Altmannshofer et al JHEP 1612 (2016) 106

- New gauge boson  $Z'$  coupling only to the **2<sup>nd</sup>** and **3<sup>rd</sup>** generation of leptons ( $L_\mu - L_\tau$ )<sup>[1]</sup> may explain:
  - long-standing  $(g-2)_\mu$  anomaly
  - dark matter abundance




- In Belle II we search for the processes:  
 $e^+e^- \rightarrow \mu^+\mu^- Z'$




# Search for an invisible $Z'$

- Search for the process:  $e^+e^- \rightarrow \mu^+\mu^-Z' \rightarrow \text{invisible}$   
→ Two possible interpretations:
  - 1) *Vanilla*,  $\text{BF}(Z' \rightarrow \nu\bar{\nu}) \sim 33\text{-}100\%$
  - 2) *Full invisible*,  $\text{BF}(Z' \rightarrow x\bar{x}) \sim 100\%$
- Look for a narrow peak **in the recoil mass against a  $\mu^+\mu^-$  pair** in events where nothing else is detected
- Dominant background radiative QED processes:
  - 1)  $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$
  - 2)  $e^+e^- \rightarrow \tau^+\tau^-(\gamma)$  (especially with both  $\tau \rightarrow \mu$ )
  - 3)  $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$
- Final State Radiation properties of the emitted  $Z'$  fed in a neural network trained for all  $Z'$  masses simultaneously





# Search for an invisible $Z'$

- The signal yield extraction is performed through a **two-dimensional fit**
  - exploit of the features in the  $M_{\text{recoil}}^2$  vs.  $\theta_{\text{recoil}}$  distribution
  - double the sensitivity with respect to the one-dimensional fit



# Search for an invisible $Z'$



- **No excess found in 79.7 fb<sup>-1</sup>**  
 → 90% CL upper limits on  $\sigma(e^+e^- \rightarrow \mu^+\mu^-Z', Z' \rightarrow \text{invisible})$  and on  $g'$   
 →  $(g-2)_\mu$  favored region excluded for  $0.8 < M(Z') < 5 \text{ GeV}/c^2$



# Search for a $\tau$ resonance in $ee \rightarrow \mu\mu\tau\tau$

- Search for a **di-tau resonance** in  $e^+e^- \rightarrow \mu^+\mu^-\tau^+\tau^-$  as a peak in the recoil against two muons
- Reconstruct  **$\tau$  decays to one-charged particle** (+ $nh^0$ )  
 → select **four-track events** with at least two tracks identified as muons  
 →  **$M(4\text{tracks}) < 9.5 \text{ GeV}/c^2$**  to suppress the four-lepton backgrounds that peak at the c.m. energy
- **Background suppression exploits features of kinematic variables in the signal** ( $X$  arising from a final state radiation, system recoiling against the 2 muons is a tau pair)

Accepted by PRL: arXiv:2306.12294






# Search for a $\tau\tau$ resonance in $ee \rightarrow \mu\mu\tau\tau$

[2] W. Altmannshofer et. al. JHEP 12 (2016) 106  
 [3] B. Batell, N. Lange, D. McKeen, M. Pospelov, and A. Ritz, Phys. Rev. D 95, 075003 (2017)  
 [4] M. Bauer, M. Neubert, and A. Thamm, J. High Energy Phys. 2017, 44 (2017)


Accepted by PRL: arXiv:2306.12294

- **No significant excess observed in  $62.8 \text{ fb}^{-1}$**   
 → 90% CL upper limits on the process cross-section  
 $\sigma(e^+e^- \rightarrow (X \rightarrow \tau^+\tau^-) \mu^+\mu^-) = \sigma(e^+e^- \rightarrow X \mu^+\mu^-) B(X \rightarrow \tau^+\tau^-)$ , with  $X = S, \text{ALP}, Z'$
- Exclusion limits on the couplings for three different models ( $Z'^{[2]}$ , **leptophilic scalar (S)**<sup>[3]</sup>, and **ALP**<sup>[4]</sup>) are derived:



# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$

- Search for the process  $e^+e^- \rightarrow \mu^+\mu^- X$ , with  $X \rightarrow \mu^+\mu^-$  ( $X = Z', S$ )  
→ Look for a peak in the opposite charge di-muon mass distribution in  $e^+e^- \rightarrow \mu^+\mu^-\mu^+\mu^-$  events
- $(L_\mu - L_\tau)$  model used as benchmark and then performances are checked for the **scalar case** <sup>[5]</sup>



- Scalar particle coupling through Yukawa-like interaction, only
- Mainly proposed as a way to solve the muon  $(g-2)_\mu$  anomaly

$$\mathcal{L} \supset (g_S S \bar{\mu} \mu)$$

Coupling constant:  
induces a shift in  
 $\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{theory}}$

- If  $m_S > 2m_\mu$  the only tree-level decay channel is  $S \rightarrow \mu\mu$   
( $S \rightarrow \nu\nu, \gamma\gamma$  also are possible at one loop level, but highly suppressed)

# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$

- Events selected have **4 charged particles**:

- zero charge
- at least **three identified as muons**
- $M(4\text{-tracks}) \sim \sqrt{s}/c^2$
- no extra energy


- Main SM background contributions:

- 1)  $e^+e^- \rightarrow \mu^+\mu^-\mu^+\mu^-$
- 2)  $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$
- 3)  $e^+e^- \rightarrow \mu^+\mu^-(\gamma)$


→ **Multi-Layer Perceptron (MLP)-based background suppression**

Signal over background discrimination relying on a few variables sensitive the signal features:



- (a) Presence of a  $\mu\mu$  resonance
- (b) Production mechanism



# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$



Peak corresponding to the **maximum muon pair momentum**





- transformed variables fed into MLP in order to reduce their change with the  $Z'$  mass
- five separate MLPs in different  $M(\mu\mu)$  intervals
- selection optimized in each interval with a figure of merit
- **background rejection factor from 2 to 14**

# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$

- **No significant excess observed in 178 fb<sup>-1</sup>**

→ 90% CL upper limits on the process cross-section  $\sigma(e^+e^- \rightarrow X \mu^+\mu^-) \times B(X \rightarrow \mu^+\mu^-)$ , with  $X = Z', S$





To be submitted to PRD

# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$

- **No significant excess observed in 178 fb<sup>-1</sup>**

→ 90% CL upper limits on the process cross-section  $\sigma(e^+e^- \rightarrow X \mu^+\mu^-) \times B(X \rightarrow \mu^+\mu^-)$ , with  $X = Z', S$

→ Cross section limits are translated into upper limits on the  $g'$  coupling constant for the  $L_\mu - L_\tau$  model and on the  $g_s$  coupling constant for the muonphilic dark scalar  $S$ <sup>[5]</sup>



To be submitted to PRD

# Conclusion

---


- Belle II/SuperKEKB is a **unique environment** to search for **light dark matter or mediators**
- **Excellent sensitivity** for dark sector searches
- **World's leading results** are obtained with a subset of the full available data
  - Search for invisible  $Z'$
  - Search for visible  $Z'$  to muons (+ muonphilic scalar)
  - Search for visible  $Z'$  to taus (+ leptophilic scalar and ALP)
- $424 \text{ fb}^{-1}$  recorded to date, **more results with higher statistics and improved analyses will be produced**

Thank you!

**Backup**



# SuperKEKB




Total beam current  
1.5 x KEKB current



$$L = \frac{\gamma_{\pm}}{2er_e} \left( \frac{I_{\pm} \xi_{y\pm}}{\beta_{y\pm}^*} \right) \left( \frac{R_L}{R_{\xi y}} \right)$$

Beta function at the IP  
20 times smaller than KEKB

# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$ : $J/\psi$




Closeup around  $J/\psi$  nominal mass



- data/MC ratio is over 1 (but for very low masses)
- Modulations due to the different MLP ranges
- Visible features:  $\rho$ ,  $J/\psi$ ,  $Y(1S)$

# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$



# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$ : muonphilic dark-scalar

We extended the  $Z'$  search to the case of a muophilic dark scalar,  $S$

- Scalar particle coupling through Yukawa-like interaction, only
- Mainly proposed as a way to solve the muon  $(g-2)_\mu$  anomaly

$$\mathcal{L} \supset g_S S \bar{\mu} \mu$$

Coupling constant:

induces a shift in

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{theory}}$$


- If  $m_S > 2m_\mu$  the only tree-level decay channel is  $S \rightarrow \mu\mu$   
( $S \rightarrow \nu\nu, \gamma\gamma$  also are possible at one loop level, but highly suppressed)

We reinterpreted our result in terms of the dark scalar  $S$ , keeping all the steps of the analysis completely unaltered

- 1) P. Harris, P. Schuster, J. Zupan, *Snowmass White Paper: New flavors and rich structures in dark sectors*
- 2) S. Gori, M. Williams, et al., *Dark Sector Physics at High-Intensity Experiments*
- 3) D. Forbes, C. Herwig, *New Searches for Muonphilic Particles at Proton Beam Dump Spectrometers*
- 4) R. Capdevilla, D. Curtin et al., *Systematically testing singlet models for  $(g-2)_\mu$*



# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$ : muonphilic dark-scalar




Difference:  $Z'$  is softly produced at low masses,  $S$  have a hard momentum spectrum also in the low mass region.

In  $e^+e^- \rightarrow \mu^+\mu^-X$  interactions  $X$  can be:

- A vector: production occurs through a s-wave process
- A scalar: production occurs through a p-wave process



**At low  $S$  masses the p-wave suppression makes the scalar process grow slowly with the energy, while there is no suppression for vector processes.**

# Search for a $\mu\mu$ resonance in $ee \rightarrow \mu\mu\mu\mu$ : muonphilic dark-scalar




# Search for a dark leptophilic scalar in $\tau$ decays at Belle

- Search for a narrow peak in  $m_{ll}$  distribution
- Mass range probed in this analysis:  
 **$40 \text{ MeV} < m(\Phi_l) < 6.5 \text{ GeV}$** 
  - $\Phi_l \rightarrow e^+e^-$  for  $m(\Phi_l) < 2m(\mu) \rightarrow$  **low mass region**
  - $\Phi_l \rightarrow \mu^+\mu^-$  for  $m(\Phi_l) > 2m(\mu) \rightarrow$  **high mass region**
- **Strategy:**
  - $\rightarrow e^+e^- \rightarrow \tau^+\tau^- \Phi_l$  require 1-prong decay
  - $\rightarrow$  4 tracks with 0 net charge
- **Background:**  $e^+e^- \rightarrow \tau^+\tau^-$ ,  $e^+e^-/\mu^+\mu^-$ ,  $q\bar{q}$ ,  $B\bar{B}$ 
  - $\rightarrow$  Define five BDT score to suppress backgrounds
- Maximum Likelihood fit to  $m_{ll}$  distribution
  - $\rightarrow$  Evaluate sensitivities to each mass point




# Search for a long-lived (pseudo-)scalar particle in $b \rightarrow s$

- Search for **dark scalar** particles  $S$  from  $B$  decays in **rare  $b \rightarrow s$  transitions**
  - $S$  could mix with SM Higgs with mixing angle  $\theta_s$  (naturally long-lived for  $\theta_s \ll 1$ )
  - $M_S < M_B$ , decays of  $S$  into dark matter particles must be kinematically forbidden to provide the correct relic density
- Look for  $S$  decays into SM final states in **8 exclusive channels**:



- **B-meson candidates** are reconstructed from prompt and displaced charged tracks
- **S candidates** are reconstructed from displaced oppositely-charged tracks pairs
- B-meson kinematics to reject combinatorial background
- **Signature:** bump hunt with extended max likelihood unbinned fits to the (\*)reduced mass spectrum, separately for each channel and lifetime





# Search for a long-lived (pseudo-)scalar particle in $b \rightarrow s$

- **No significant excess found in 189 fb<sup>-1</sup>**
  - first model-independent 95% CL upper limits on **BF(B→KS)×BF(S→x<sup>+</sup>x<sup>-</sup>)**
  - translate into model independent limits on  $\sin\theta_s$  vs.  $m_s$

First limits  
on decay  
to hadrons



Results are  
also available  
for the  
pseudo-scalar  
(ALP) model



Submitted to PRL: <https://arxiv.org/abs/2306.02830>

# Search for a dark leptophilic scalar in $\tau$ decays at Belle

- No significant excess observed in  $626 \text{ fb}^{-1}$  in all mass region



- 90 % CL UL on  $\xi$  vs  $m(\Phi_L)$ 
  - Comparable or more stringent limits than BaBar (Phys. Rev. Lett. 125, 181801)
  - Exclude a wide range of parameter space of the model favored by  $(g-2)_\mu$

